

# A LOCAL CRITERION FOR WEYL MODULES FOR GROUPS OF TYPE A

VLADIMIR SHCHIGOLEV

**ABSTRACT.** Let  $G$  be a universal Chevalley group over an algebraically closed field and  $\mathcal{U}^-$  be the subalgebra of  $\text{Dist}(G)$  generated by all divided powers  $X_{\alpha,m}$  with  $\alpha < 0$ . We conjecture an algorithm to determine if  $Fe_{\omega}^+ \neq 0$ , where  $F \in \mathcal{U}^-$ ,  $\omega$  is a dominant weight and  $e_{\omega}^+$  is a highest weight vector of the Weyl module  $\Delta(\omega)$ . This algorithm does not use bases of  $\Delta(\omega)$  and is similar to the algorithm for irreducible modules that involves stepwise raising the vector under investigation. For an arbitrary  $G$ , this conjecture is proved in one direction and for  $G$  of type A in both.

## 1. INTRODUCTION

Let  $\Sigma$  be a root system and  $\mathbf{K}$  be an algebraically closed field. Consider the semisimple complex Lie algebra  $\mathfrak{L}$  with Cartan subalgebra  $\mathcal{H}$  and root system  $\Sigma$ . For each  $\alpha \in \Sigma$ , we denote by  $H_{\alpha}$  the element of  $\mathcal{H}$  introduced in [4, Lemma 1].

We denote by  $\Sigma^+$  a positive root system of  $\Sigma$  and by  $\Pi = \{\alpha_1, \dots, \alpha_{\ell}\}$  the simple root system contained in  $\Sigma^+$ . We choose elements  $X_{\alpha}$ , where  $\alpha \in \Sigma$ , so that they together with  $H_{\alpha_1}, \dots, H_{\alpha_{\ell}}$  form a Chevalley basis of  $\mathfrak{L}$  in the sense of [4, Theorem 1]. Let  $G$  be the universal Chevalley group over  $\mathbf{K}$  constructed by this basis as in [4, § 3].

Following [4, Theorem 2], we denote by  $\mathcal{U}_{\mathbf{Z}}$  the subring of the universal enveloping algebra of  $\mathfrak{L}$  generated by  $X_{\alpha}^m/m!$ , where  $\alpha \in \Sigma$  and  $m \in \mathbf{Z}^+$  (the set of nonnegative integers). In this paper, we shall consider modules over the algebra  $\mathcal{U} := \mathcal{U}_{\mathbf{Z}} \otimes_{\mathbf{Z}} \mathbf{K}$ , which is called the *hyperalgebra* of  $G$  (denoted by  $\text{Dist}(G)$  in [2]). As a  $\mathbf{K}$ -algebra  $\mathcal{U}$  is generated by the elements  $X_{\alpha,m} := (X_{\alpha}^m/m!) \otimes 1_{\mathbf{K}}$ . Moreover, every rational  $G$ -module  $V$  can be made into a  $\mathcal{U}$ -module by the rule

$$x_{\alpha}(t)v = \sum_{m=0}^{+\infty} t^m X_{\alpha,m}v,$$

where  $v \in V$ ,  $\alpha \in \Sigma$ ,  $t \in \mathbf{K}$  and  $x_{\alpha}(t)$  is the root element of  $G$  corresponding to  $\alpha$  and  $t$  (see [4, § 3]). We shall also need the elements  $H_{\alpha,m} = \binom{H_{\alpha}}{m} \otimes 1_{\mathbf{K}}$ . It is easy to show that these elements actually belong to  $\mathcal{U}$  (e.g. [4, Corollary to Lemma 5]). The reader should keep in mind the formula  $\binom{H_{\alpha}+r}{m} \otimes 1_{\mathbf{K}} = \sum_{n=0}^m H_{\alpha,n} \binom{r}{m-n}$ , where  $r \in \mathbf{Z}$ .

**Proposition 1.** *The products  $\prod_{\alpha \in \Sigma^+} X_{-\alpha, m_{-\alpha}} \cdot \prod_{i=1}^{\ell} H_{\alpha_i, n_i} \cdot \prod_{\alpha \in \Sigma^+} X_{\alpha, m_{\alpha}}$ , where  $m_{-\alpha}, n_i, m_{\alpha} \in \mathbf{Z}^+$ , taken in some fixed order form a basis of  $\mathcal{U}$ .*

We shall always mean the grading of  $\mathcal{U}$  in which  $X_{\alpha,m}$  has weight  $m\alpha$  and  $H_{\alpha,m}$  has weight 0. Elements of  $\mathcal{U}$  that are homogeneous with respect to this grading will be called *weight elements*.

We denote the  $\mathbf{K}$ -span of all the above basis elements

- with unitary second and third factors by  $\mathcal{U}^-$ ;
- with unitary first and third factors by  $\mathcal{U}^0$ ;
- with unitary third factor by  $\mathcal{U}^{-,0}$ .

It can be easily checked that each of these spaces is a  $\mathbf{K}$ -algebra. A vector  $v$  of a rational  $G$ -module is called *primitive* if  $X_{\alpha,m}v = 0$  for any  $\alpha \in \Sigma^+$  and  $m > 0$  and is called *simply primitive* if  $X_{\alpha,1}v = 0$  for any  $\alpha \in \Sigma^+$ .

To the Chevalley basis of  $\mathfrak{L}$ , we associate the maximal torus  $T$  in the standard way (see [4, Lemma 21], where the torus is denoted by  $H$ ). We denote by  $X(T)$  and  $X^+(T)$  the set of weights and the set of dominant (with respect to  $\Sigma^+$ ) weights of  $T$  respectively. Let  $\omega_1, \dots, \omega_\ell$  be the fundamental weights of  $T$  corresponding to the roots  $\alpha_1, \dots, \alpha_\ell$ . Each weight  $\omega \in X(T)$  has the form  $\omega = a_1\omega_1 + \dots + a_\ell\omega_\ell$ , where  $a_i \in \mathbf{Z}$ . Under this notation, we set  $h(\omega) := a_1 + \dots + a_\ell$ . For each  $\omega \in X^+(T)$ , we denote by  $\Delta(\omega)$  and  $\nabla(\omega)$  the Weyl module and the co-Weyl module with highest weight  $\omega$  respectively. We also fix some nonzero vector  $e_\omega^+$  of  $\Delta(\omega)$  having weight  $\omega$ . In what follows, we denote by  $X_{\alpha,m}^V$  the operator on a rational  $G$ -module  $V$  acting as the left multiplication by  $X_{\alpha,m}$ . We also denote by  $V^\tau$  for  $\tau \in X(T)$  the  $\tau$ -weight space of  $V$ . A *weigh decomposition* of an element  $x$  belonging to  $\mathcal{U}$  or to a rational  $G$ -module is  $x = x_1 + \dots + x_m$ , where  $x_1, \dots, x_m$  are weight elements of mutually distinct weights.

It is well known that a vector of the irreducible rational  $G$ -module  $L(\omega)$  is nonzero if and only if it can be raised up to proportionality to the highest weight vector  $v_\omega^+$  of  $L(\omega)$  by successive multiplications by the elements  $X_{\alpha,m}$  for  $\alpha \in \Sigma^+$ . Thus for any  $F \in \mathcal{U}^-$ , we can determine whether  $Fv_\omega^+ \neq 0$  by calculating directly in  $\mathcal{U}$  (see Remark 4). Note that we should at first consider the weight decomposition  $F = F_1 + \dots + F_m$  and then investigate if  $F_i v_\omega^+ \neq 0$  separately for each  $i$ .

Conjectures A and B in Section 2 suggest algorithms similar to this algorithm that concern the Weyl module  $\Delta(\omega)$  instead of the irreducible module  $L(\omega)$ . These conjectures are proved in one direction, see Corollary 7. This result may be useful for establishing nonzero homomorphism between Weyl modules. Simple examples are given in Sections 2.5 and 2.6. In these examples, we choose  $Y$  to be the corresponding elements described at page 241 of [5] evaluated at  $\omega$ . The coefficients in the commutator relations for  $B_2$  can be found in [8, §25 Exercise 6].

Conjectures A and B turn out to hold for  $G = A_\ell(\mathbf{K})$ . This is proved in Section 3. We use there the standard basis theorem for Weyl modules, the straightening rule and the results of Section 3.1 concerning some properties of graphs of special type called flows.

Section 4 contains the list of notations globally used throughout this paper.

**Acknowledgments.** The author would like to thank Irina Suprunenko for drawing his attention to this problem and a useful discussion.

## 2. CONJECTURES AND REFORMULATIONS

**2.1. Conjectures.** Given a weight  $\omega \in X(T)$ , we define the  $\mathbf{K}$ -linear map  $\text{ev}^\omega : \mathcal{U}^{-,0} \rightarrow \mathcal{U}^-$  by

$$\text{ev}^\omega \left( \prod_{\alpha \in \Sigma^+} X_{-\alpha, m_{-\alpha}} \cdot \prod_{i=1}^\ell H_{\alpha_i, n_i} \right) = \prod_{\alpha \in \Sigma^+} X_{-\alpha, m_{-\alpha}} \cdot \prod_{i=1}^\ell \binom{\omega(H_{\alpha_i})}{n_i}.$$

The map  $\text{ev}^\omega$  enjoys the properties

**(ev-1)** the restriction  $\text{ev}^\omega|_{\mathcal{U}^0}$  is a  $\mathbf{K}$ -algebra homomorphism from  $\mathcal{U}^0$  to  $\mathbf{K}$ ;

**(ev-2)** if  $v$  is a vector of weight  $\omega$  of a rational  $G$ -module and  $P \in \mathcal{U}^{-,0}$ , then  $Pv = \text{ev}^\omega(P)v$ .

Given a weight  $\omega \in X(T)$ , a root  $\alpha \in \Sigma^+$  and an integer  $m \in \mathbf{Z}^+$ , we define the operator  $r_{\alpha, m}^\omega$  on  $\mathcal{U}^-$  as follows. Let  $F$  belong to  $\mathcal{U}^-$ . By Proposition 1, we have the unique representation  $X_{\alpha, m}F = P + E$ , where  $P \in \mathcal{U}^{-,0}$  and  $E$  belongs to the left ideal of  $\mathcal{U}$  generated by the elements  $X_{\beta, k}$  with  $\beta \in \Sigma^+$  and  $k > 0$ . Then we set  $r_{\alpha, m}^\omega(F) := \text{ev}^\omega(P)$ . The map  $r_{\alpha, m}^\omega$  enjoys the properties

**(r-1)** if  $F$  has weight  $\tau$  then  $r_{\alpha, m}^\omega(F)$  has weight  $\tau + m\alpha$ ;

**(r-2)** if  $v$  is a primitive vector of weight  $\omega$  of a rational  $G$ -module,  $F \in \mathcal{U}^-$ ,  $\alpha \in \Sigma^+$  and  $m \in \mathbf{Z}^+$ , then  $X_{\alpha, m}Fv = r_{\alpha, m}^\omega(F)v$ ;

*Example.* We have  $X_{\alpha_i, 1}X_{-\alpha_i, 2} = X_{-\alpha_i, 1}(H_{\alpha_i, 1} - 1) + X_{-\alpha_i, 2}X_{\alpha_i, 1}$ . Therefore,  $r_{\alpha_i, 1}^\omega(X_{-\alpha_i, 2}) = \text{ev}^\omega(X_{-\alpha_i, 1}(H_{\alpha_i, 1} - 1)) = (a_i - 1)X_{-\alpha_i, 1}$ , where  $\omega = a_1\omega_1 + \dots + a_\ell\omega_\ell$ .

To formulate our conjectures, we consider the following transformations on  $\mathcal{U}^- \times X^+(T)$ :

- (a)  $(F, \omega) \mapsto (r_{\alpha, m}^\omega(F), \omega)$ , where  $\alpha \in \Sigma^+$  and  $m \in \mathbf{Z}^+$ ;
- (b)  $(F, \omega) \mapsto (F, \omega - \delta)$ , where  $\delta$  is a weight of  $X^+(T)$  such that  $\omega - \delta \in X^+(T)$ .

**Definition 2.** Let  $F \in \mathcal{U}^-$  and  $\omega \in X^+(T)$ . The pair  $(F, \omega)$  is called *reducible* (simply reducible) to a pair  $(c, 0)$ , where  $c \in \mathbf{K}^*$ , if there exists a sequence

$$(F^{(k)}, \omega^{(k)}), \dots, (F^{(0)}, \omega^{(0)}) \quad (1)$$

of pairs of  $\mathcal{U}^- \times X^+(T)$  such that

- (i)  $(F^{(k)}, \omega^{(k)}) = (F, \omega)$  and  $(F^{(0)}, \omega^{(0)}) = (c, 0)$ ;
- (ii) for any  $i = 0, \dots, k-1$ ,  $(F^{(i)}, \omega^{(i)})$  is derived from  $(F^{(i+1)}, \omega^{(i+1)})$  by transformation (a) (resp. transformation (a) with  $m = 1$ ) or transformation (b).

**Remark 3.** If  $(F, \omega)$  is simply reducible to  $(c, 0)$ , then  $(F, \omega)$  is reducible to  $(c, 0)$ .

We shall also say that a pair  $(F, \omega)$  is *reducible* (simply reducible) if this pair is reducible (simply reducible) to some pair  $(c, 0)$ , where  $c \in \mathbf{K}^*$ .

**Conjecture A.** Let  $F$  be a weight element of  $\mathcal{U}^-$  and  $\omega \in X^+(T)$ . We have  $Fe_\omega^+ \neq 0$  if and only if the pair  $(F, \omega)$  is reducible.

**Conjecture B.** Let  $F$  be a weight element of  $\mathcal{U}^-$  and  $\omega \in X^+(T)$ . We have  $Fe_\omega^+ \neq 0$  if and only if the pair  $(F, \omega)$  is simply reducible.

**Remark 4.** Obviously, for a weight element  $F$  of  $\mathcal{U}^-$ ,  $Fv_\omega^+ \neq 0$  if and only if  $F$  can be reduced to  $c \in \mathbf{K}^*$  by the transformations  $F \mapsto r_{\alpha,m}^\omega(F)$ .

*Example.* The Weyl module  $\Delta(0)$  is one-dimensional and therefore is irreducible. Take  $F = 1 + X_{-\alpha}$ , where  $\alpha \in \Sigma^+$ . Then  $Fe_0^+ = e_0^+ \neq 0$ , while the pair  $(F, 0)$  is not reducible.

The above example shows why we consider only weight elements in Conjectures A and B. However, each of these conjectures if proved would allow us to answer whether  $Fe_\omega^+ \neq 0$  for an arbitrary  $F \in \mathcal{U}^-$ . Indeed, consider the weight decomposition  $F = F_1 + \cdots + F_m$ . Then  $Fe_\omega^+ \neq 0$  if and only if  $F_i e_\omega^+ \neq 0$  for some  $i$ .

**2.2. Proof in one direction.** Of principal importance for the theory developed in this paper is the following result.

**Lemma 5.** *Let  $\omega, \delta \in X^+(T)$  such that  $\omega - \delta \in X^+(T)$ . Then there exists the  $\mathcal{U}^-$ -homomorphism  $d_\delta^\omega : \Delta(\omega) \rightarrow \Delta(\omega - \delta)$  that takes  $e_\omega^+$  to  $e_{\omega-\delta}^+$ .*

*Proof.* Let  $w_0$  be the longest element of the Weyl group of  $\Sigma$ . Suppose temporarily that  $\text{char } \mathbf{K} = 0$ . Then  $L(\delta)^* \cong L(-w_0\delta)$  by [2, Corollary II.2.5 and Proposition II.2.6]. Since  $v_{\omega-\delta}^+ \otimes v_\delta^+$  is a nonzero primitive vector of  $L(\omega - \delta) \otimes L(\delta)$  having weight  $\omega$ , the universal property of Weyl modules [2, Lemma II.2.13 b] implies the existence of a nonzero homomorphism from  $\Delta(\omega)$  to  $L(\omega - \delta) \otimes L(\delta)$ . Using  $\Delta(\omega) \cong L(\omega)$  due to  $\text{char } \mathbf{K} = 0$  and [2, Lemma I.4.4], we obtain

$$\begin{aligned} K &\cong \text{Hom}_G(L(\omega), L(\omega - \delta) \otimes L(\delta)) \cong \text{Hom}_G(L(\omega) \otimes L(\delta)^*, L(\omega - \delta)) \\ &\cong \text{Hom}_G(L(\omega) \otimes L(-w_0\delta), L(\omega - \delta)). \end{aligned}$$

Hence  $L(\omega - \delta)$  is a composition factor of  $L(\omega) \otimes L(-w_0\delta)$  with multiplicity 1. Here we used that any rational  $G$ -module is semisimple. Let  $L(\tau)$  be a composition factor of  $L(\omega) \otimes L(-w_0\delta)$  with  $\tau \neq \omega - \delta$ . Again using the above mentioned fact on semisimplicity, we get

$$\begin{aligned} 0 &\neq \text{Hom}_G(L(\omega) \otimes L(-w_0\delta), L(\tau)) \cong \text{Hom}_G(L(\omega), L(\tau) \otimes L(-w_0\delta)^*) \\ &= \text{Hom}_G(L(\omega), L(\tau) \otimes L(\delta)), \end{aligned}$$

whence  $\omega \leq \tau + \delta$  and  $\omega - \delta < \tau$ . In terms of characters, these facts can be written as

$$\text{ch } L(\omega) \otimes L(-w_0\delta) = \text{ch } L(\omega - \delta) + \sum_{\tau > \omega - \delta} a_\tau \text{ch } L(\tau) \quad (2)$$

for some  $a_\tau \in \mathbf{Z}^+$ .

Now let us return to the situation where  $\mathbf{K}$  has arbitrary characteristic. Then (2) can be rewritten as

$$\text{ch } \nabla(\omega) \otimes \nabla(-w_0\delta) = \text{ch } \nabla(\omega - \delta) + \sum_{\tau > \omega - \delta} a_\tau \text{ch } \nabla(\tau). \quad (3)$$

The main result of [3] implies that  $\nabla(\omega) \otimes \nabla(-w_0\delta)$  has a good filtration. The factors of this filtration are given by (3). Therefore applying [2, II.4.16 Remark 4], we obtain that  $\nabla(\omega - \delta)$  is the bottom factor in some good filtration of  $\nabla(\omega) \otimes \nabla(-w_0\delta)$ . The module  $\Delta(\omega) \otimes \Delta(-w_0\delta)$  is contravariantly dual to  $\nabla(\omega) \otimes \nabla(-w_0\delta)$ , whence it has  $\Delta(\omega - \delta)$  as a top factor.

Let  $\pi : \Delta(\omega) \otimes \Delta(-w_0\delta) \rightarrow \Delta(\omega - \delta)$  be the corresponding epimorphism of  $G$ -modules.

Let  $u$  be a nonzero vector of  $\Delta(-w_0\delta)$  of minimal weight  $-\delta$ . We are going to prove that  $e_\omega^+ \otimes u \notin \ker \pi$ . Suppose on the contrary that  $e_\omega^+ \otimes u \in \ker \pi$ . The comparison of weights shows that  $e_\omega^+ \otimes (\bigoplus_{\tau > -\delta} \Delta(-w_0\delta)^\tau) \subset \ker \pi$ . Since  $\dim \Delta(-w_0\delta)^{-\delta} = \dim \Delta(-w_0\delta)^{-w_0\delta} = 1$ , we obtain  $e_\omega^+ \otimes \Delta(-w_0\delta) \subset \ker \pi$ .

It is a well-known fact that  $e_\omega^+ \otimes \Delta(-w_0\delta)$  generates  $\Delta(\omega) \otimes \Delta(-w_0\delta)$  as a  $G$ -module. We sketch the proof for completeness. Let  $W$  denote the  $G$ -submodule of  $\Delta(\omega) \otimes \Delta(-w_0\delta)$  generated by  $e_\omega^+ \otimes \Delta(-w_0\delta)$ . Suppose that  $W \neq \Delta(\omega) \otimes \Delta(-w_0\delta)$  and take  $\sigma_{\max}$  to be a maximal weight such that  $\Delta(\omega)^{\sigma_{\max}} \otimes \Delta(-w_0\delta)$  is not contained in  $W$ . Let  $v_1 \in \Delta(\omega)^{\sigma_{\max}}$  and  $v_2 \in \Delta(-w_0\delta)$  be arbitrary vectors. We have  $v_1 = Fe_\omega^+$  for some  $F \in \mathcal{U}^-$  having weight  $\sigma_{\max} - \omega$ . Obviously,  $W \ni F(e_\omega^+ \otimes v_2) = v_1 \otimes v_2 + w$ , where  $w \in \bigoplus_{\sigma > \sigma_{\max}} \Delta(\omega)^\sigma \otimes \Delta(-w_0\delta)$ . The latter sum is contained in  $W$  by the choice of  $\sigma_{\max}$ , whence  $w \in W$  and  $v_1 \otimes v_2 \in W$ . We proved  $\Delta(\omega)^{\sigma_{\max}} \otimes \Delta(-w_0\delta) \subset W$ , which is a contradiction.

Hence  $\ker \pi = \Delta(\gamma) \otimes \Delta(-w_0\delta)$ , since  $\ker \pi$  is a  $G$ -submodule. This contradicts  $\pi \neq 0$ . We proved that  $e_\omega^+ \otimes u \notin \ker \pi$ . Hence  $\pi(e_\omega^+ \otimes u) = ce_{\omega-\delta}^+$ , where  $c \in \mathbf{K}^*$ . Now the required map is given by  $d_\delta^\omega(v) = c^{-1}\pi(v \otimes u)$ , where  $v \in \Delta(\omega)$ .  $\square$

We note the following trivial property of these homomorphisms:

**(d-1)** if  $\omega, \tau, \omega - \delta, \delta - \tau \in X^+(T)$  then  $d_\delta^\omega = d_{\delta-\tau}^{\omega-\tau} d_\tau^\omega$ .

**Lemma 6.**  $\ker d_\delta^\omega$  is a  $T$ -module.

*Proof.* Let  $v \in \ker d_\delta^\omega$ . Consider the weight decomposition  $v = v_1 + \dots + v_m$ , where each  $v_i \neq 0$ . It suffices to show that each  $v_i \in \ker d_\delta^\omega$ .

There exist weight elements  $F_1, \dots, F_m$  of  $\mathcal{U}^-$  such that  $v_i = F_i e_\omega^+$  for any  $i = 1, \dots, m$ . We have

$$0 = d_\delta^\omega(v) = d_\delta^\omega(v_1) + \dots + d_\delta^\omega(v_m) = F_1 e_{\omega-\delta}^+ + \dots + F_m e_{\omega-\delta}^+.$$

Since the weights of  $F_1, \dots, F_m$  are mutually distinct, we obtain  $0 = F_i e_{\omega-\delta}^+ = d_\delta^\omega(v_i)$  for any  $i = 1, \dots, m$ .  $\square$

**Corollary 7.** Let  $F \in \mathcal{U}^-$  and  $\omega \in X^+(T)$ . If  $(F, \omega)$  is reducible then  $Fe_\omega^+ \neq 0$ .

*Proof.* We apply induction on the length  $k+1$  of sequence (1) satisfying properties (i) and (ii) of Definition 2. For  $k=0$ , the result is obvious, since in that case  $F = c \in \mathbf{K}^*$ .

Now suppose that  $k > 0$ . Then the inductive hypothesis yields

$$F^{(k-1)} e_{\omega^{(k-1)}}^+ \neq 0, \quad (4)$$

since the sequence  $(F^{(k-1)}, \omega^{(k-1)}), \dots, (c, 0)$  has length  $k$  and we assume that for sequences of length smaller than  $k+1$  the result is true.

*Case 1:*  $(F^{(k-1)}, \omega^{(k-1)})$  is derived from  $(F, \omega)$  by transformation (a). In that case, we have  $\omega^{(k-1)} = \omega$  and  $F^{(k-1)} = r_{\alpha, m}^\omega(F)$  for some  $\alpha \in \Sigma^+$  and  $m \in \mathbf{Z}^+$ . By (r-2) and (4) we obtain

$$X_{\alpha, m} F e_\omega^+ = r_{\alpha, m}^\omega(F) e_\omega^+ = F^{(k-1)} e_{\omega^{(k-1)}}^+ \neq 0,$$

whence  $Fe_\omega^+ \neq 0$ .

*Case 2:*  $(F^{(k-1)}, \omega^{(k-1)})$  is derived from  $(F, \omega)$  by transformation (b). In that case, we have  $F^{(k-1)} = F$  and  $\omega^{(k-1)} = \omega - \delta$  for some  $\delta \in X^+(T)$ . By Lemma 5 and (4) we obtain

$$d_\delta^\omega(Fe_\omega^+) = Fd_\delta^\omega(e_\omega^+) = Fe_{\omega-\delta}^+ = F^{(k-1)}e_{\omega^{(k-1)}}^+ \neq 0,$$

whence again  $Fe_\omega^+ \neq 0$ .  $\square$

It follows from this corollary and Remark 3 that Conjecture B implies Conjecture A as follows:

$$(F, \omega) \text{ is reducible} \xrightarrow{\text{Cor.7}} Fe_\omega^+ \neq 0,$$

$$Fe^+ \neq 0 \xrightarrow{\text{Con.B}} (F, \omega) \text{ is simply reducible} \xrightarrow{\text{Rem.3}} (F, \omega) \text{ is reducible.}$$

**2.3. Reformulations.** The following result will be used in Section 3.

**Lemma 8.** *Conjecture B holds for all  $(F, \omega)$  if and only if for any nonzero  $\omega \in X^+(T)$ , the intersection*

$$\bigcap \{ \ker d_{\omega_i}^\omega \mid i = 1, \dots, \ell \text{ and } \omega - \omega_i \in X^+(T) \} \quad (5)$$

*does not contain nonzero simply primitive vectors.*

*Proof.* “Only if” part. Suppose that  $\omega$  is a nonzero dominant weight and (5) contains a nonzero simply primitive vector  $v$ . By Lemma 6, we can assume that  $v$  is a weight vector. We obviously have  $v = Fe_\omega^+$  for some weight element  $F$  of  $\mathcal{U}^-$ . We claim that  $(F, \omega)$  is not simply reducible, i.e. Conjecture B is violated for  $(F, \omega)$ .

Suppose, on the contrary, that  $(F, \omega)$  is simply reducible to  $(c, 0)$ , where  $c \in \mathbf{K}^*$ . Then there exists a sequence of form (1) satisfying properties (i) and (ii) of Definition 2, where in (ii)  $m$  always equals 1 if transformation (a) is applied. Without loss of generality we may suppose that in each transformation (b) used to derive  $(F^{(i)}, \omega^{(i)})$  from  $(F^{(i+1)}, \omega^{(i+1)})$ , we take  $\delta \neq 0$ . We have  $k > 0$ , since otherwise  $F = c$  and  $v \notin \ker d_{\omega_i}^\omega$  for any  $i = 1, \dots, \ell$  such that  $\omega - \omega_i \in X^+(T)$ . Note that at least one such  $i$  exists since  $\omega \neq 0$ . Hence  $v$  does not belong to (5).

Like any pair of sequence (1), the pair  $(F^{(k-1)}, \omega^{(k-1)})$  is reducible. Thus Corollary 7 implies

$$F^{(k-1)}e_{\omega^{(k-1)}}^+ \neq 0. \quad (6)$$

*Case 1:*  $(F^{(k-1)}, \omega^{(k-1)})$  is derived from  $(F, \omega)$  by transformation (a) with  $m = 1$ . In that case, we have  $\omega^{(k-1)} = \omega$  and  $F^{(k-1)} = r_{\alpha,1}^\omega(F)$  for some  $\alpha \in \Sigma^+$ . Applying the fact that  $v$  is simply primitive and (r-2), we obtain

$$0 = X_{\alpha,1}v = X_{\alpha,1}Fe_\omega^+ = r_{\alpha,1}^\omega(F)e_\omega^+ = F^{(k-1)}e_{\omega^{(k-1)}}^+,$$

which contradicts (6).

*Case 2:*  $(F^{(k-1)}, \omega^{(k-1)})$  is derived from  $(F, \omega)$  by transformation (b). In that case, we have  $F^{(k-1)} = F$  and  $\omega^{(k-1)} = \omega - \delta$  for some nonzero weight  $\delta$  of  $X^+(T)$ . Since we supposed that  $\delta \neq 0$ , there exists  $j = 1, \dots, \ell$  such that  $\delta - \omega_j \in X^+(T)$ . Since  $\omega^{(k-1)} \in X^+(T)$ , we clearly have  $\omega - \omega_j =$

$\omega^{(k-1)} + \delta - \omega_j \in X^+(T)$ . Thus  $v \in \ker d_{\omega_j}^\omega$ , since  $v$  belongs to (5). By (d-1), we obtain

$$0 = d_{\delta - \omega_j}^{\omega - \omega_j} d_{\omega_j}^\omega(v) = d_\delta^\omega(v) = F d_\delta^\omega(e_\omega^+) = F e_{\omega - \delta}^+ = F^{(k-1)} e_{\omega^{(k-1)}}^+,$$

which again contradicts (6).

*“If” part.* Suppose that for any nonzero dominant weight  $\omega$ , intersection (5) does not contain nonzero simply primitive vectors. In view of Corollary 7, we must only prove that for any  $F \in \mathcal{U}^-$  having weight  $\tau$  and  $\omega \in X^+(T)$  such that  $F e_\omega^+ \neq 0$ , the pair  $(F, \omega)$  is simply reducible.

We apply induction on  $(-\tau, h(\omega))$  with componentwise order on such pairs. If  $\omega = 0$  then  $F e_\omega^+ \neq 0$  implies that  $\tau = 0$  and  $F$  is a nonzero element of  $\mathbf{K}$ . In that case  $(F, \omega)$  is simply reducible (to itself). Therefore, we suppose that  $\omega \neq 0$ .

*Case 1:  $F e_\omega^+$  is not simply primitive.* We have  $X_{\alpha,1} F e_\omega^+ \neq 0$  for some  $\alpha \in \Sigma^+$ . By (r-2), we obtain

$$0 \neq X_{\alpha,1} F e_\omega^+ = r_{\alpha,1}^\omega(F) e_\omega^+.$$

By (r-1), the vector  $r_{\alpha,1}^\omega(F)$  has weight  $\tau + \alpha \leq 0$ . Since  $(-(\tau + \alpha), h(\omega))$  is smaller than  $(-\tau, h(\omega))$ , the inductive hypothesis implies that  $(r_{\alpha,1}^\omega(F), \omega)$  is simply reducible. It remains to notice that  $(r_{\alpha,1}^\omega(F), \omega)$  is derived from  $(F, \omega)$  by transformation (a) with  $m = 1$ .

*Case 2:  $F e_\omega^+$  does not belong to (5).* There exists some  $j = 1, \dots, \ell$  such that  $\omega - \omega_j \in X^+(T)$  and  $F e_\omega^+ \notin \ker d_{\omega_j}^\omega$ . We have

$$0 \neq d_{\omega_j}^\omega(F e_\omega^+) = F d_{\omega_j}^\omega(e_\omega^+) = F e_{\omega - \omega_j}^+.$$

Since  $h(\omega - \omega_j) = h(\omega) - 1$ , the pair  $(-\tau, h(\omega - \omega_j))$  is smaller than the pair  $(-\tau, h(\omega))$  and the inductive hypothesis implies that  $(F, \omega - \omega_j)$  is simply reducible. It remains to notice that  $(F, \omega - \omega_j)$  is derived from  $(F, \omega)$  by transformation (b).

It is natural to consider the remaining case when we are in neither Case 1 nor Case 2. Then  $F e_\omega^+$  is a nonzero simply primitive vector belonging to (5). Since  $\omega \neq 0$ , this contradicts the assumption of this part of the proof.  $\square$

Arguing as above but dropping the word “simply” and the condition  $m=1$ , we obtain the following result.

**Lemma 9.** *Conjecture A holds if and only if for any nonzero  $\omega \in X^+(T)$ , intersection (5) does not contain nonzero primitive vectors (equivalently, nonzero  $G$ -submodules).*

**2.4. Twisting operators.** Now we shift attention to the hyperalgebra by introducing for any  $\delta \in X(T)$  the *twisting operator*  $\theta_\delta$  on  $\mathcal{U}^{-,0}$  as the  $\mathbf{K}$ -linear map defined by

$$\theta_\delta \left( \prod_{\alpha \in \Sigma^+} X_{-\alpha, m_{-\alpha}} \cdot \prod_{i=1}^\ell H_{\alpha_i, n_i} \right) = \prod_{\alpha \in \Sigma^+} X_{-\alpha, m_{-\alpha}} \cdot \prod_{i=1}^\ell \left( H_{\alpha_i + \delta(H_{\alpha_i})}^{n_i} \right) \otimes 1_{\mathbf{K}}.$$

The map  $\theta_\delta$  enjoys the properties

( $\theta$ -1)  $\theta_\delta$  is a  $\mathbf{K}$ -algebra automorphism of  $\mathcal{U}^{-,0}$ ;

( $\theta$ -2) if  $\omega, \delta, \omega - \delta \in X^+(T)$ ,  $P \in \mathcal{U}^{-,0}$  and  $v \in \Delta(\omega)$ , then

$$d_\delta^\omega(Pv) = \theta_\delta(P)d_\delta^\omega(v).$$

We need to comment only on ( $\theta$ -2). By linearity, it is enough to consider only the case  $P = F'H$ , where  $F' \in \mathcal{U}^-$  and  $H \in \mathcal{U}^0$ , and  $v = Fe_\omega^+$ , where  $F$  is an element of  $\mathcal{U}^-$  having weight  $\tau$ . Applying (ev-2), ( $\theta$ -1), the definition of  $\theta_\delta$  and the fact that  $d_\delta^\omega$  is a  $\mathcal{U}^-$ -homomorphism, we obtain

$$\begin{aligned} d_\delta^\omega(Pv) &= F'd_\delta^\omega(Hv) = F'd_\delta^\omega(\text{ev}^{\omega+\tau}(H)Fe_\omega^+) \\ &= F'\text{ev}^{\omega+\tau}(H)Fd_\delta^\omega(e_\omega^+) = \text{ev}^{\omega+\tau}(H)F'Fe_{\omega-\delta}^+, \\ \theta_\delta(P)d_\delta^\omega(v) &= F'\theta_\delta(H)Fd_\delta^\omega(e_\omega^+) = F'\theta_\delta(H)Fe_{\omega-\delta}^+ \\ &= \text{ev}^{\omega-\delta+\tau}(\theta_\delta(H))F'Fe_{\omega-\delta}^+. \end{aligned}$$

Hence it remains to prove that  $\text{ev}^{\omega-\delta+\tau}(\theta_\delta(H)) = \text{ev}^{\omega+\tau}(H)$ . By (ev-1) and ( $\theta$ -1), it is enough to consider the case  $H = H_{\alpha_i, n}$ . We have

$$\begin{aligned} \text{ev}^{\omega-\delta+\tau}(\theta_\delta(H_{\alpha_i, n})) &= \text{ev}^{\omega-\delta+\tau}\left(\left(H_{\alpha_i} + \delta(H_{\alpha_i})\right)_n \otimes 1_{\mathbf{K}}\right) = \\ &= \left((\omega-\delta+\tau)(H_{\alpha_i}) + \delta(H_{\alpha_i})\right)_n \otimes 1_{\mathbf{K}} = \left((\omega+\tau)(H_{\alpha_i})\right)_n \otimes 1_{\mathbf{K}} = \text{ev}^{\omega+\tau}(H_{\alpha_i, n}). \end{aligned}$$

For any  $\alpha \in \Sigma^+$  and  $m \in \mathbf{Z}^+$ , we define the  $\mathbf{K}$ -linear operator  $\eta_{\alpha, m}$  on  $\mathcal{U}^{-,0}$  as follows. Let  $F$  belong to  $\mathcal{U}^-$ . Then  $X_{\alpha, m}F = P + E$ , where  $P \in \mathcal{U}^{-,0}$  and  $E$  belongs to the left ideal of  $\mathcal{U}$  generated by the elements  $X_{\beta, k}$  with  $\beta \in \Sigma^+$  and  $k > 0$ . We set  $\eta_{\alpha, m}(F) := P$ . The last element already was introduced, when we defined  $r_{\alpha, m}^\omega$ . We clearly have  $\text{ev}^\omega(\eta_{\alpha, m}(F)) = r_{\alpha, m}^\omega(F)$ . The operator  $\eta_{\alpha, m}$  enjoys the property

( $\eta$ -1) if  $v$  is a primitive vector of a rational  $G$ -module,  $F \in \mathcal{U}^-$ ,  $\alpha \in \Sigma^+$  and  $m \in \mathbf{Z}^+$ , then  $X_{\alpha, m}Fv = \eta_{\alpha, m}(F)v$ .

**Lemma 10.** *Let  $\Phi(x_1, \dots, x_k, y)$  be an element of the free associative algebra over  $\mathbf{K}$  with generators  $x_1, \dots, x_k, y$  (noncommutative associative polynomial) linear in  $y$  and  $\omega, \delta, \omega - \delta \in X^+(T)$ . Let  $D_\delta^\omega$  be any operator on the module  $V := \Delta(\omega) \oplus \Delta(\omega - \delta)$  whose restriction to  $\Delta(\omega)$  coincides with  $d_\delta^\omega$ . We have*

$$\Phi(X_{\beta_1, m_1}^V, \dots, X_{\beta_k, m_k}^V, D_\delta^\omega)(Fe_\omega^+) = \Phi(\eta_{\beta_1, m_1}, \dots, \eta_{\beta_k, m_k}, \theta_\delta)(F) \cdot e_{\omega-\delta}^+$$

for any  $\beta_1, \dots, \beta_k \in \Sigma^+$ ,  $m_1, \dots, m_k \in \mathbf{Z}^+$  and  $F \in \mathcal{U}^-$ .

*Proof.* By linearity, it suffices to prove that

$$\begin{aligned} X_{\beta_1, m_1}^V \cdots X_{\beta_q, m_q}^V D_\delta^\omega X_{\beta_{q+1}, m_{q+1}}^V \cdots X_{\beta_k, m_k}^V (Fe_\omega^+) \\ = \eta_{\beta_1, m_1} \cdots \eta_{\beta_q, m_q} \theta_\delta \eta_{\beta_{q+1}, m_{q+1}} \cdots \eta_{\beta_k, m_k} (F) \cdot e_{\omega-\delta}^+. \end{aligned} \quad (7)$$

By ( $\eta$ -1) and ( $\theta$ -2), the left-hand side of (7) equals

$$\begin{aligned} X_{\beta_1, m_1} \cdots X_{\beta_q, m_q} d_\delta^\omega (X_{\beta_{q+1}, m_{q+1}} \cdots X_{\beta_k, m_k} Fe_\omega^+) \\ = X_{\beta_1, m_1} \cdots X_{\beta_q, m_q} d_\delta^\omega \left( \eta_{\beta_{q+1}, m_{q+1}} \cdots \eta_{\beta_k, m_k} (F) \cdot e_\omega^+ \right) \\ = X_{\beta_1, m_1} \cdots X_{\beta_q, m_q} \left( \theta_\delta \eta_{\beta_{q+1}, m_{q+1}} \cdots \eta_{\beta_k, m_k} (F) \right) \cdot e_{\omega-\delta}^+ \\ = \eta_{\beta_1, m_1} \cdots \eta_{\beta_q, m_q} \theta_\delta \eta_{\beta_{q+1}, m_{q+1}} \cdots \eta_{\beta_k, m_k} (F) \cdot e_{\omega-\delta}^+, \end{aligned}$$



which is exactly the right-hand side of (7).  $\square$

**2.5. Example for  $G = B_2(\mathbf{K})$  and root  $2\alpha + \beta$ .** We are going to show a possible application of Corollary 7. Let  $\text{char } \mathbf{K} = p > 2$ ,  $G = B_2(\mathbf{K})$  and  $\Sigma^+ = \{\alpha, \beta\}$ , where  $\alpha$  is short and  $\beta$  is long. We denote by  $\omega_\alpha$  and  $\omega_\beta$  the fundamental weights corresponding to  $\alpha$  and  $\beta$  respectively. Take a dominant weight  $\omega = a\omega_\alpha + b\omega_\beta$  such that  $a \geq 2$ ,  $b \geq 1$  and  $a + b + 1 \equiv 0 \pmod{p}$ . We claim that  $\text{Hom}_G(\Delta(\omega - (2\alpha + \beta)), \Delta(\omega)) \neq 0$ . Indeed, consider the elements

$$\begin{aligned} Y &= X_{-\alpha,1}^2 X_{-\beta,1} - N_{\alpha,\beta} b X_{-\alpha,1} X_{-\alpha-\beta,1} + b(b+1) \frac{N_{\alpha,\beta} N_{\alpha,\alpha+\beta}}{2} X_{-2\alpha-\beta,1}, \\ Z &= 2X_{-\alpha,1} X_{-\beta,1} - N_{\alpha,\beta} b X_{-\alpha-\beta,1}, \end{aligned}$$

where  $[X_\delta, X_\gamma] = N_{\delta,\gamma} X_{\delta+\gamma}$  for  $\delta, \gamma, \delta+\gamma \in \Sigma$ . Elementary calculations show

$$r_{\alpha,1}^\omega(Y) = (a+b+1)Z, \quad r_{\alpha,2}^\omega(Y) = (a+b+2)(a+b+1)X_{-\beta,1}, \quad r_{\beta,1}^\omega(Y) = 0.$$

The equivalence  $a + b + 1 \equiv 0 \pmod{p}$  and (r-2) imply that  $Ye_\omega^+$  is a primitive vector of  $\Delta(\omega)$  having weight  $\omega - (2\alpha + \beta)$ . We are going to show that this vector is nonzero by showing that the pair  $(Y, \omega)$  is reducible. Choose any integers  $a'$  and  $b'$  such that  $0 \leq a' \leq a$  and  $0 \leq b' \leq b$  and set  $\omega' := a'\omega_\alpha + b'\omega_\beta$ . Consider the following sequence of transformations:

$$\begin{aligned} (Y, \omega) &\xrightarrow[\text{transf. (b)}]{\delta=\omega'} (Y, \omega-\omega') \xrightarrow[\text{transf. (a)}]{r_{\alpha,1}^{\omega-\omega'}} ((a-a'+b+1)Z, \omega-\omega') \xrightarrow[\text{transf. (a)}]{r_{\beta,1}^{\omega-\omega'}} \\ &(a'(2b'-b)X_{-\alpha,1}, \omega-\omega') \xrightarrow[\text{transf. (a)}]{r_{\alpha,1}^{\omega-\omega'}} (a'(2b'-b)(a-a')1_{\mathbf{K}}, \omega-\omega') \xrightarrow[\text{transf. (b)}]{\delta=\omega-\omega'} \\ &(a'(2b'-b)(a-a')1_{\mathbf{K}}, 0). \end{aligned}$$

To ensure that  $a'(2b'-b)(a-a') \not\equiv 0 \pmod{p}$ , we set  $a' := 1$  if  $a \not\equiv 1 \pmod{p}$ ,  $a' := 2$  if  $a \equiv 1 \pmod{p}$ ,  $b' := 0$  if  $b \not\equiv 0 \pmod{p}$  and  $b' := 1$  if  $b \equiv 0 \pmod{p}$ .

**2.6. Example for  $G = B_2(\mathbf{K})$  and root  $\alpha + \beta$ .** In this example, we use the same notation as in the previous one but assume that  $\text{char } \mathbf{K}$  is an arbitrary prime  $p$ . Take a dominant weight  $\omega = a\omega_\alpha + b\omega_\beta$  such that  $a \geq 1$ ,  $b \geq 1$  and  $a + 2b + 2 \equiv 0 \pmod{p}$ . We claim that  $\text{Hom}_G(\Delta(\omega - (\alpha + \beta)), \Delta(\omega)) \neq 0$ . Indeed, consider the element

$$Y = X_{-\alpha,1} X_{-\beta,1} - N_{\alpha,\beta} b X_{-\alpha-\beta,1}.$$

Elementary calculations show

$$r_{\alpha,1}^\omega(Y) = (a + 2b + 2)X_{-\beta,1}, \quad r_{\beta,1}^\omega(Y) = 0.$$

The equivalence  $a + 2b + 2 \equiv 0 \pmod{p}$  and (r-2) imply that  $Ye_\omega^+$  is a primitive vector of  $\Delta(\omega)$  having weight  $\omega - (\alpha + \beta)$ . We are going to show that this vector is nonzero by showing that the pair  $(Y, \omega)$  is reducible.

Choose any integers  $a'$  and  $b'$  such that  $0 \leq a' \leq a$  and  $0 \leq b' \leq b$  and set  $\omega' := a'\omega_\alpha + b'\omega_\beta$ . Consider the following sequence of transformations:

$$\begin{aligned} (Y, \omega) &\xrightarrow[\text{transf. (b)}]{\delta=\omega'} (Y, \omega-\omega') \xrightarrow[\text{transf. (a)}]{r_{\alpha,1}^{\omega-\omega'}} ((a-a'+2b+2)X_{-\beta,1}, \omega-\omega') \xrightarrow[\text{transf. (a)}]{r_{\beta,1}^{\omega-\omega'}} \\ &(a'(b'-b)1_{\mathbf{K}}, \omega-\omega') \xrightarrow[\text{transf. (b)}]{\delta=\omega-\omega'} (a'(b'-b)1_{\mathbf{K}}, 0). \end{aligned}$$

To ensure that  $a'(b'-b) \not\equiv 0 \pmod{p}$ , we set  $a' := 1$  and  $b' := 0$  if  $b \not\equiv 0 \pmod{p}$  and  $b' := 1$  if  $b \equiv 0 \pmod{p}$ .

### 3. CRITERION FOR $G = A_\ell(\mathbf{K})$ .

**3.1. Flows.** In what follows, we call  $s$  the *beginning* and  $t$  the *end* of  $(s, t)$  or say that  $(s, t)$  *begins* at  $s$  and *ends* at  $t$ .

**Definition 11.** A directed graph  $\Gamma$  is called a *flow* with sources  $a_1, \dots, a_q$  and sinks  $b_1, \dots, b_q$  if

- (i) the set of vertices of  $\Gamma$  is  $\mathbf{Z}$ ; every edge of  $\Gamma$  has the form  $(s, t)$  with  $s < t$ ; the number of edges of  $\Gamma$  is finite;
- (ii) for each  $j = 1, \dots, q$ , there is exactly one edge of  $\Gamma$  beginning at  $a_j$ ;
- (iii) for each  $j = 1, \dots, q$ , there is exactly one edge of  $\Gamma$  ending at  $b_j$ ;
- (iv) for each integer  $r$  distinct from  $a_1, \dots, a_q, b_1, \dots, b_q$ , either no edge of  $\Gamma$  ends or begins at  $r$  or exactly one edge of  $\Gamma$  ends at  $r$  and exactly one edge of  $\Gamma$  begins at  $r$ . In the latter case,  $r$  is called a *transit point* of  $\Gamma$ .

For  $q = 0$ , the only possible flow is the *empty flow*  $\Gamma_\emptyset$ , which has no edges.

We denote by  $S_q$  the symmetric group thought of as the group of all bijections of  $\{1, \dots, q\}$ . We assume that the elements of  $S_q$  act on this set on the left and multiply them accordingly. We also identify  $S_q$  with a subgroup of  $S_{q+1}$  assuming that  $\sigma(q+1) = q+1$  for any  $\sigma \in S_q$ .

We denote the set of all flows  $\Gamma$  with sources  $a_1, \dots, a_q \leq i$  and sinks  $b_1, \dots, b_q > i$  having no transit point greater than  $i$  by  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ . Clearly,  $\Gamma$  decomposes into connected components. If there is a component of  $\Gamma$  containing  $a_j$  and  $b_k$ , then we say that source  $a_j$  and sink  $b_k$  are linked in  $\Gamma$ . Throughout this section, we suppose for definiteness that  $a_1 < a_2 < \dots < a_q$  and  $b_q < b_{q-1} < \dots < b_1$ . Let  $\sigma$  denote the permutation in  $S_q$  such that  $a_j$  and  $b_{\sigma(j)}$  are linked in  $\Gamma$ . We shall occasionally call  $\sigma$  the *linking permutation* of  $\Gamma$ . We define the *i-sign* of  $\Gamma$  to be

$$\text{sgn}_i(\Gamma) := \text{sgn } \sigma \cdot (-1)^{\sum_{j=1}^q (b_j - i) + \text{number of transit points of } \Gamma}. \quad (8)$$

*Example.* Let  $\Gamma$  have the following edges  $(5, 9), (3, 8), (6, 7), (4, 6), (2, 4), (1, 3)$ . Then  $\Gamma \in \mathcal{F}_6(1, 2, 5; 9, 8, 7)$  and has transit points 3, 4, 6. Its linking permutation is  $\sigma$  with  $\sigma(1) = 2, \sigma(2) = 3$  and  $\sigma(3) = 1$ . Therefore, its 6-sign is  $-1$ .

We define the operators  $L_1, L_2, L_3, M_1, M_2, R$  on some flows  $\Gamma$  of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  as follows.

| Graph         | Condition                                                              | Operation                                        |
|---------------|------------------------------------------------------------------------|--------------------------------------------------|
| $L_1(\Gamma)$ | $q \geq 1$ ,<br>$a_q - 1$ is a transit point of $\Gamma$               | replacement of $(s, a_q - 1)$<br>with $(s, a_q)$ |
| $L_2(\Gamma)$ | $q \geq 1$ ,<br>no edge of $\Gamma$ begins at $a_q - 1$                | addition of $(a_q - 1, a_q)$                     |
| $L_3(\Gamma)$ | $q \geq 1$ ,<br>$a_q - 1$ is no source of $\Gamma$                     | replacement of $(a_q, t)$<br>with $(a_q - 1, t)$ |
| $M_1(\Gamma)$ | $i + 1$ is no sink of $\Gamma$ ,<br>$i$ is a transit point of $\Gamma$ | replacement of $(s, i)$<br>with $(s, i + 1)$     |
| $M_2(\Gamma)$ | $i + 1$ is no sink of $\Gamma$ ,<br>no edge of $\Gamma$ begins at $i$  | addition of $(i, i + 1)$                         |
| $R(\Gamma)$   | $q \geq 1$ ,<br>$b_q + 1$ is no sink of $\Gamma$                       | replacement of $(s, b_q)$<br>with $(s, b_q + 1)$ |

Table 1

In each row of this table, the graph in the left column is obtained from  $\Gamma$  by the operation described in the right column (here  $s$  and  $t$  can be chosen uniquely) if  $\Gamma$  satisfies the condition in the middle column.

*Example.* Let  $\Gamma$  be as in the previous example. Then  $L_1(\Gamma)$  has edges  $(5, 9)$ ,  $(3, 8)$ ,  $(6, 7)$ ,  $(4, 6)$ ,  $(2, 5)$ ,  $(1, 3)$ ;  $L_2(\Gamma)$  is not well-defined, since  $a_q - 1 = a_3 - 1 = 4$  is a transit point of  $\Gamma$ ;  $L_3(\Gamma)$  has edges  $(4, 9)$ ,  $(3, 8)$ ,  $(6, 7)$ ,  $(4, 6)$ ,  $(2, 4)$ ,  $(1, 3)$  and is no flow (see Lemma 14).

**Lemma 12.** *If well-defined,  $L_1(\Gamma)$  belongs to  $\mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$  and has  $i$ -sign opposite to that of  $\Gamma$ .*

*Proof.* In  $L_1(\Gamma)$ ,  $a_q$  becomes a transit point and  $a_q - 1$  becomes a source. Therefore,  $L_1(\Gamma)$  and  $\Gamma$  have the same sinks and number of transit points.

Let  $\sigma$  be the linking permutation of  $\Gamma$  and  $a_l$  be the source belonging to the connected component of  $\Gamma$  containing  $a_q - 1$ . Clearly,  $l < q$ .

We set  $a'_j := a_j$  if  $j = 1, \dots, q - 1$  and  $a'_q := a_q - 1$ . Then  $a'_1, \dots, a'_q$  are the sources of  $L_1(\Gamma)$  arranged in the strictly ascending order. We set  $\tau := \sigma \circ (l, q)$ . One can easily see that  $a'_j$  and  $b_{\tau(j)}$  are linked in  $L_1(\Gamma)$  for any  $j = 1, \dots, q$ , i.e.  $\tau$  is the linking permutation of  $L_1(\Gamma)$ .  $\square$

**Lemma 13.** *If well-defined,  $L_2(\Gamma)$  belongs to  $\mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$  and has  $i$ -sign opposite to that of  $\Gamma$ .*

*Proof.* In  $L_2(\Gamma)$ ,  $a_q$  becomes a transit point and  $a_q - 1$  becomes a source. Therefore,  $L_2(\Gamma)$  has the same sinks as  $\Gamma$  and one more transit point than  $\Gamma$ . Finally notice that  $L_2(\Gamma)$  and  $\Gamma$  have the same linking permutations.  $\square$

**Lemma 14.** *Suppose that  $L_3(\Gamma)$  is well-defined. Then  $L_3(\Gamma)$  is a flow if and only if  $a_q - 1$  is no transit point of  $\Gamma$ . If  $L_3(\Gamma)$  is a flow, then it belongs to  $\mathcal{F}_i(a_1, \dots, a_{q-1}; a_q - 1, b_1, \dots, b_q)$  and has the same  $i$ -sign as  $\Gamma$ . If  $L_3(\Gamma)$  is no flow, then there is exactly one flow  $\Gamma'$  of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  distinct from  $\Gamma$  such that  $L_3(\Gamma) = L_3(\Gamma')$ . Moreover,  $\text{sgn}_i(\Gamma) = -\text{sgn}_i(\Gamma')$ .*

*Proof.* If  $a_q - 1$  is no transit point, then  $L_3(\Gamma)$  is a flow with the same linking permutation, sinks and number of transit points. Hence  $\text{sgn}_i(L_3(\Gamma)) = \text{sgn}_i(\Gamma)$ .

Now consider the case where  $a_q - 1$  is a transit point of  $\Gamma$ . Then there is an edge of  $\Gamma$  of the form  $(a_q - 1, r)$ . On the other hand, there is also an edge of  $\Gamma$  of the form  $(a_q, t)$ , since  $a_q$  is a source of  $\Gamma$ . In  $L_3(\Gamma)$ ,  $(a_q, t)$  turns to  $(a_q - 1, t)$ , whence  $L_3(\Gamma)$  has two edges  $(a_q - 1, t)$  and  $(a_q - 1, r)$  beginning at  $a_q - 1$  and  $L_3(\Gamma)$  is no flow.

Consider the flow  $\Gamma'$  that is obtained from  $\Gamma$  by the removing the edges  $(a_q - 1, r)$  and  $(a_q, t)$  and adding the edges  $(a_q, r)$  and  $(a_q - 1, t)$ . It is straightforward to see that  $\Gamma'$  is the required flow. To calculate the  $i$ -sign of  $\Gamma'$ , denote by  $l$  the integer such that  $a_l$  belongs to the connected component of  $\Gamma$  containing  $r$  and denote by  $\sigma$  the linking permutation of  $\Gamma$ . Clearly,  $l < q$ . Similarly to Lemma 13, one can see that  $\sigma \circ (l, q)$  is the linking permutation of  $\Gamma'$ . Moreover,  $\Gamma$  and  $\Gamma'$  have the same transit points and sinks.  $\square$

**Lemma 15.** *If well-defined,  $M_1(\Gamma)$  belongs to  $\mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i+1)$  and has  $i$ -sign opposite to that of  $\Gamma$ .*

*Proof.* For  $M_1(\Gamma)$ ,  $i$  is no longer a transit point (but a source) and  $i + 1$  is a new sink. Therefore the second factor of the right-hand side of (8) does not change.

Let  $\sigma$  be the linking permutation of  $\Gamma$  and  $a_l$  be the source belonging to the connected component of  $\Gamma$  containing  $i$ . Note that  $a_1, \dots, a_q, i$  are all the sources of  $M_1(\Gamma)$  written in the strictly ascending order and  $b_1, \dots, b_q, i + 1$  are all the sinks of  $M_1(\Gamma)$  written in the strictly descending order. Therefore, under the natural identification  $S_q < S_{q+1}$ , we obtain that  $\sigma \circ (l, q + 1)$  is the linking permutation of  $M_1(\Gamma)$ .  $\square$

**Lemma 16.** *If well-defined,  $M_2(\Gamma)$  belongs to  $\mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i+1)$  and has  $i$ -sign opposite to that of  $\Gamma$ .*

*Proof.* Note that  $a_1, \dots, a_q, i$  are all the sources of  $M_2(\Gamma)$  written in the strictly ascending order and  $b_1, \dots, b_q, i + 1$  are all the sinks of  $M_2(\Gamma)$  written in the strictly descending order. Hence the linking permutation of  $M_2(\Gamma)$  is the same as that of  $\Gamma$  under the natural identification  $S_q < S_{q+1}$ . Moreover,  $M_2(\Gamma)$  has the same transit points as  $\Gamma$ .  $\square$

**Lemma 17.** *If well-defined,  $R(\Gamma)$  belongs to  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_{q-1}, b_q + 1)$  and has  $i$ -sign opposite to that of  $\Gamma$ .*

*Proof.* Note that  $R(\Gamma)$  has the same transit points and sources as  $\Gamma$ . Moreover,  $b_1, \dots, b_{q-1}, b_q + 1$  are all the sinks of  $M_2(\Gamma)$  written in the strictly descending order. In particular,  $R(\Gamma)$  and  $\Gamma$  have the same linking permutation.  $\square$

Now we are going to define “inverse maps” for each group of operators  $L_1, L_2, L_3; M_1, M_2; R$ .

Suppose that  $q \geq 1$ ,  $a_q - 1 \notin \{a_1, \dots, a_{q-1}\}$  and  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$ . We define the operator  $\mathcal{L}^o(\Gamma')$  that is one of the operators  $L_1, L_2, L_3$  described in Table 1 and the graph  $\mathcal{L}^g(\Gamma')$  using the table below.

| Condition                                                      |                          | Value                                                                  |
|----------------------------------------------------------------|--------------------------|------------------------------------------------------------------------|
| there is an edge $(s, a_q)$ of $\Gamma'$<br>with $s < a_q - 1$ | $\mathcal{L}^o(\Gamma')$ | $L_1$                                                                  |
|                                                                | $\mathcal{L}^g(\Gamma')$ | obtained from $\Gamma'$ by replacing<br>$(s, a_q)$ with $(s, a_q - 1)$ |
| $(a_q - 1, a_q)$ is an edge of $\Gamma'$                       | $\mathcal{L}^o(\Gamma')$ | $L_2$                                                                  |
|                                                                | $\mathcal{L}^g(\Gamma')$ | obtained from $\Gamma'$ by<br>removing $(a_q - 1, a_q)$                |
| $a_q$ is no transit point of $\Gamma'$                         | $\mathcal{L}^o(\Gamma')$ | $L_3$                                                                  |
|                                                                | $\mathcal{L}^g(\Gamma')$ | obtained from $\Gamma'$ by replacing<br>$(a_q - 1, t)$ with $(a_q, t)$ |

Table 2

Let us take any row of this table such that the condition in its left column is satisfied. If the middle column of this row contains  $\mathcal{L}^g(\Gamma')$ , then this graph is obtained from the flow  $\Gamma'$  by the operation described in the right column. If the middle column contains  $\mathcal{L}^o(\Gamma')$ , then this operator equals the operator in the right column.

*Example.* Let  $q = 2$ ,  $i = 3$ ,  $a_1 = 1$ ,  $a_2 = 3$ ,  $b_2 = 4$ ,  $b_1 = 5$  and  $\Gamma'$  has edges  $(2, 5)$ ,  $(3, 4)$ ,  $(1, 3)$ . Then  $\mathcal{L}^o(\Gamma') = L_1$  and  $\mathcal{L}^g(\Gamma')$  has edges  $(2, 5)$ ,  $(3, 4)$ ,  $(1, 2)$  (two top rows of Table 2 were applied). If we apply  $\mathcal{L}^o(\Gamma')$  to  $\mathcal{L}^g(\Gamma')$ , then we obtain  $\Gamma'$  (see the definition of  $L_1$  in Table 1).

**Lemma 18.** *Suppose that  $q \geq 1$  and  $a_q - 1 \notin \{a_1, \dots, a_{q-1}\}$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$ , we have  $\mathcal{L}^g(\Gamma') \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and  $\mathcal{L}^o(\Gamma')(\mathcal{L}^g(\Gamma')) = \Gamma'$ . For any flow  $\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and integer  $k = 1, 2, 3$ , we have  $\mathcal{L}^o(L_k(\Gamma)) = L_k$  and  $\mathcal{L}^g(L_k(\Gamma)) = \Gamma$  if  $L_k(\Gamma)$  is well-defined and a flow.*

*Proof.* The result follows directly from Tables 1 and 2.  $\square$

**Corollary 19.** *Suppose that  $q \geq 1$  and  $a_q - 1 \notin \{a_1, \dots, a_{q-1}\}$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$ , there exists a unique pair  $(O, \Gamma) \in \{L_1, L_2, L_3\} \times \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $O(\Gamma) = \Gamma'$ .*

*Proof.* It follows from the first assertion of Lemma 18 that  $(\mathcal{L}^o(\Gamma'), \mathcal{L}^g(\Gamma'))$  is a suitable pair. Now suppose that  $(L_{k_1}, \Gamma_1)$  and  $(L_{k_2}, \Gamma_2)$  are pairs of  $\{L_1, L_2, L_3\} \times \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $L_{k_1}(\Gamma_1) = L_{k_2}(\Gamma_2) = \Gamma'$ . By the second assertion of Lemma 18, we have

$$\begin{aligned} L_{k_1} &= \mathcal{L}^o(L_{k_1}(\Gamma_1)) = \mathcal{L}^o(L_{k_2}(\Gamma_2)) = L_{k_2}, \\ \Gamma_1 &= \mathcal{L}^g(L_{k_1}(\Gamma_1)) = \mathcal{L}^g(L_{k_2}(\Gamma_2)) = \Gamma_2. \end{aligned}$$

$\square$

Now suppose that  $a_q < i$ ,  $i + 1 < b_q$  if  $q \geq 1$  and  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i + 1)$ . Then we use the table below to define the operator  $\mathcal{M}^o(\Gamma')$  and the graph  $\mathcal{M}^g(\Gamma')$ .

| Condition                                             |                          | Value                                                         |
|-------------------------------------------------------|--------------------------|---------------------------------------------------------------|
| there is an edge $(s, i+1)$ of $\Gamma'$ with $s < i$ | $\mathcal{M}^o(\Gamma')$ | $M_1$                                                         |
|                                                       | $\mathcal{M}^g(\Gamma')$ | obtained from $\Gamma'$ by replacing $(s, i+1)$ with $(s, i)$ |
| $(i, i+1)$ is an edge of $\Gamma'$                    | $\mathcal{M}^o(\Gamma')$ | $M_2$                                                         |
|                                                       | $\mathcal{M}^g(\Gamma')$ | obtained from $\Gamma'$ by removing $(i, i+1)$                |

Table 3

This table defines the new graphs and the new operators in the same way as Table 2.

**Lemma 20.** *Suppose that  $a_q < i$ ,  $i+1 < b_q$  if  $q \geq 1$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i+1)$ , we have  $\mathcal{M}^g(\Gamma') \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and  $\mathcal{M}^o(\Gamma')(\mathcal{M}^g(\Gamma')) = \Gamma'$ . For any flow  $\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and integer  $k = 1, 2$ , we have  $\mathcal{M}^o(M_k(\Gamma)) = M_k$  and  $\mathcal{M}^g(M_k(\Gamma)) = \Gamma$  if  $M_k(\Gamma)$  is well-defined.*

*Proof.* The result follows directly from Tables 1 and 3.  $\square$

**Corollary 21.** *Suppose that  $a_q < i$ ,  $i+1 < b_q$  if  $q \geq 1$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i+1)$ , there exists a unique pair  $(O, \Gamma) \in \{M_1, M_2\} \times \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $O(\Gamma) = \Gamma'$ .*

*Proof.* The result follows from Lemma 20 similarly to Corollary 19.  $\square$

Finally suppose that  $q \geq 1$ ,  $b_q + 1 \notin \{b_{q-1}, \dots, b_1\}$  and  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_{q-1}, b_q + 1)$ . Then we denote by  $\mathcal{R}^g(\Gamma')$  the flow obtained from  $\Gamma'$  by replacing the edge  $(s, b_q + 1)$  with  $(s, b_q)$ .

**Lemma 22.** *Suppose that  $q \geq 1$  and  $b_q + 1 \notin \{b_{q-1}, \dots, b_1\}$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_{q-1}, b_q + 1)$ , we have  $\mathcal{R}^g(\Gamma') \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and  $R(\mathcal{R}^g(\Gamma')) = \Gamma'$ . For any flow  $\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ , we have  $\mathcal{R}^g(R(\Gamma)) = \Gamma$ .*

**Corollary 23.** *Suppose that  $q \geq 1$  and  $b_q + 1 \notin \{b_{q-1}, \dots, b_1\}$ . For any flow  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_{q-1}, b_q + 1)$ , there exists a unique flow  $\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $R(\Gamma) = \Gamma'$ .*

*Proof.* The result follows from Lemma 22 similarly to Corollary 19.  $\square$

**3.2. Basis of the hyperalgebra.** In the remainder of the paper, we consider the case  $G = A_\ell(\mathbf{K})$ . We set  $n := \ell + 1$ , order the simple roots  $\alpha_1, \dots, \alpha_{n-1}$  so that they form the diagram

$$\begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & \alpha_3 & & & & \alpha_{n-2} & & \alpha_{n-1} \end{array}$$

and use the short hand notation

$$[i_1, \dots, i_k]_i := [\eta_{\alpha_{i_1}}, \dots, \eta_{\alpha_{i_k}}, \theta_{\omega_i}],$$

where  $i_1, \dots, i_k, i$  belong to  $\{1, \dots, n-1\}$ . Here and in what follows,  $\eta_\alpha := \eta_{\alpha,1}$ ,  $[x, y] = xy - yx$  and long commutators are right-normed, i. e. are defined by the inductive rule  $[x_1, \dots, x_k] = [x_1, [x_2, \dots, x_k]]$  for  $k > 2$ .

We denote by  $\lambda_i$  the  $i$ th entry of a sequence  $\lambda$  and set  $\delta_{\mathcal{P}} := 1$  if  $\mathcal{P}$  is true and  $\delta_{\mathcal{P}} := 0$  if  $\mathcal{P}$  is false for an arbitrary condition  $\mathcal{P}$ .

In the present case, we can take  $\mathcal{L} = \mathfrak{sl}_n(\mathbf{C})$  with the standard choice of the Chevalley basis:

$$X_{\alpha_i + \dots + \alpha_{j-1}} = E_{i,j}, \quad X_{-\alpha_i - \dots - \alpha_{j-1}} = E_{j,i}, \quad H_{\alpha_i} = E_{i,i} - E_{i+1,i+1},$$

where  $1 \leq i < j \leq n$ . Here and in what follows,  $E_{i,j}$  denotes the  $n \times n$  matrix with 1 in the intersection of row  $i$  and column  $j$  and 0 elsewhere.

We denote  $E_{i,j}^{(m)} := (E_{i,j}^m / m!) \otimes 1_{\mathbf{K}}$  for  $m \in \mathbf{Z}^+$  and  $H_i := H_{\alpha_i} \otimes 1_{\mathbf{K}}$ . This is simply more convenient notation for the elements introduced in Section 1, in the case  $G = A_\ell(\mathbf{K})$ . Indeed, for  $1 \leq i < j \leq n$  we have

$$E_{i,j}^{(m)} = X_{\alpha_i + \dots + \alpha_{j-1}, m}, \quad E_{j,i}^{(m)} = X_{-\alpha_i - \dots - \alpha_{j-1}, m}, \quad H_i = H_{\alpha_i, 1}.$$

We also denote  $E_{i,j}^{(1)}$  by  $E_{i,j}$ . This notation does not cause confusion, since we shall work only in  $\mathcal{U}$ .

We extend  $E_{i,j}^{(m)}$  to negative values of  $m$  by setting  $E_{i,j}^{(m)} := 0$  if  $m < 0$ . This definition, useless at first sight, is in fact very significant, in particular, because the formulas

$$\begin{aligned} [E_{i,j}, E_{j,k}^{(m)}] &= E_{i,k} E_{j,k}^{(m-1)}, \quad [E_{i,j}, E_{k,i}^{(m)}] = -E_{k,j} E_{k,i}^{(m-1)}, \\ [E_{i,i+1}, E_{i+1,i}^{(m)}] &= E_{i+1,i}^{(m-1)} (H_i + 1 - m), \\ H_l E_{i,j}^{(m)} &= E_{i,j}^{(m)} (H_l + m(\delta_{l=i} - \delta_{l=j} - \delta_{l+1=i} + \delta_{l+1=j})), \\ E_{i,j}^{(m)} E_{i,j} &= (m+1) E_{i,j}^{(m+1)} \end{aligned} \tag{9}$$

hold for any  $m \in \mathbf{Z}$  and mutually distinct  $i, j, k = 1, \dots, n$ . These formulas are trivial for  $m < 0$  and can be proved by induction on  $m$  for  $m \geq 0$ .

More generally, the first three formulas of (9) are a special case of formulas (2.1) from [6], which also hold for any integer values of the superscripts. The reader is referred to [7] to see why, for example, setting  $\binom{x}{m} = 0$  for  $m < 0$  helps simplify a lot of calculations. The same thing happens here: the proof of Lemma 24 would be much longer if we considered  $E_{i,j}^{(m)}$  only for  $m \geq 0$ .

**3.3. Action of  $\eta_{\alpha_i}$ .** Let  $\text{UT}(n)$  denote the set of  $n \times n$  matrices  $N$  with entries in  $\mathbf{Z}$  such that  $N_{a,b} = 0$  unless  $a < b$ . We denote by  $N^s$  the sum of elements in row  $s$  of  $N$ , i.e.  $N^s := \sum_{b=1}^n N_{s,b}$ . For any  $N \in \text{UT}(n)$ , we set

$$F^{(N)} := \prod_{1 \leq a < b \leq n} E_{b,a}^{(N_{a,b})},$$

where  $E_{b,a}^{(N_{a,b})}$  precedes  $E_{d,c}^{(N_{c,d})}$  if and only if  $b < d$  or  $b = d$  and  $a < c$ . It follows from Proposition 1 that any element of  $\mathcal{U}^{-,0}$  is representable as  $\sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N$ , where  $\mathcal{H}_N \in \mathcal{U}^0$ .

**Lemma 24.** *Let  $l = 1, \dots, n-1$  and  $\mathcal{H}_N$ , where  $N \in \text{UT}(n)$ , be elements of  $\mathcal{U}^0$  that are nonzero only for finitely many matrices  $N$ . We have*

$$\eta_{\alpha_l} \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \sum_{N \in \text{UT}(n)} F^{(N)} \left( \sum_{1 \leq s < l} N_{s,l} \mathcal{H}_{N+E_{s,l+1}-E_{s,l}} \right. \\ \left. + (H_l - N^l + N^{l+1}) \mathcal{H}_{N+E_{l,l+1}} - \sum_{l+1 < t \leq n} N_{l+1,t} \mathcal{H}_{N+E_{l,t}-E_{l+1,t}} \right). \quad (10)$$

*Proof.* First we prove that

$$[E_{l,l+1}, F^{(N)}] = \sum_{1 \leq s < l} (N_{s,l} + 1) F^{(N-E_{s,l+1}+E_{s,l})} + \\ F^{(N-E_{l,l+1})} (H_l + 1 - N^l + N^{l+1}) - \sum_{l+1 < t \leq n} (N_{l+1,t} + 1) F^{(N-E_{l,t}+E_{l+1,t})} \quad (11)$$

for any  $N \in \text{UT}(n)$ . Consider the representation  $F^{(N)} = F_2 \cdots F_n$ , where  $F_j = E_{j,1}^{(N_{1,j})} \cdots E_{j,j-2}^{(N_{j-2,j})} E_{j,j-1}^{(N_{j-1,j})}$ . Since  $E_{l,l+1}$  commutes with  $F_2, \dots, F_l$ , we have  $E_{l,l+1} F^{(N)} = F_2 \cdots F_l E_{l,l+1} F_{l+1} \cdots F_n$ . Applying (9), we obtain

$$E_{l,l+1} F_{l+1} = \sum_{1 \leq s < l} \left( E_{l,s} \prod_{1 \leq r \leq l} E_{l+1,r}^{(N_{r,l+1}-\delta_{r=s})} \right) + \\ \left( \prod_{1 \leq r < l} E_{l+1,r}^{(N_{r,l+1})} \right) E_{l,l+1} E_{l,l+1}^{(N_{l,l+1})}. \quad (12)$$

Applying (9) again, we obtain

$$E_{l,l+1} E_{l+1,l}^{(N_{l,l+1})} F_{l+2} \cdots F_n = E_{l+1,l}^{(N_{l,l+1})} E_{l,l+1} F_{l+2} \cdots F_n \\ + E_{l+1,l}^{(N_{l,l+1}-1)} (H_l + 1 - N_{l,l+1}) F_{l+2} \cdots F_n = E_{l+1,l}^{(N_{l,l+1})} E_{l,l+1} F_{l+2} \cdots F_n \\ + E_{l+1,l}^{(N_{l,l+1}-1)} F_{l+2} (H_l + 1 - N_{l,l+1} - N_{l,l+2} + N_{l+1,l+2}) F_{l+3} \cdots F_n = \cdots \\ = E_{l+1,l}^{(N_{l,l+1})} E_{l,l+1} F_{l+2} \cdots F_n + E_{l+1,l}^{(N_{l,l+1}-1)} F_{l+2} \cdots F_n (H_l + 1 - N^l + N^{l+1}).$$

Thus, multiplying (12) by  $F_2 \cdots F_l$  on the left and by  $F_{l+2} \cdots F_n$  on the right, we obtain

$$E_{l,l+1} F^{(N)} = \sum_{1 \leq s < l} \left( F_2 \cdots F_l E_{l,s} \prod_{1 \leq r \leq l} E_{l+1,r}^{(N_{r,l+1}-\delta_{r=s})} \right) \times \\ \times F_{l+2} \cdots F_n + F_2 \cdots F_l \left( \prod_{1 \leq r < l} E_{l+1,r}^{(N_{r,l+1})} \right) E_{l+1,l}^{(N_{l,l+1}-1)} \times \\ \times F_{l+2} \cdots F_n (H_l + 1 - N^l + N^{l+1}) + F_2 \cdots F_{l+1} E_{l,l+1} F_{l+2} \cdots F_n = \\ \sum_{1 \leq s < l} (N_{s,l} + 1) F^{(N-E_{s,l+1}+E_{s,l})} + F^{(N-E_{l,l+1})} (H_l + 1 - N^l + N^{l+1}) + \\ F_2 \cdots F_{l+1} E_{l,l+1} F_{l+2} \cdots F_n.$$

In the above rearrangement, we used that

$$F_l E_{l,s} = (N_{s,l} + 1) \prod_{1 \leq r < l} E_{l,r}^{(N_{r,l}+\delta_{r=s})} \text{ for } 1 \leq s < l.$$



It remains to calculate  $E_{l,l+1}F_{l+2}\cdots F_n$ . Let  $t = l+2, \dots, n$ . All factors of  $F_t$  commute with  $E_{l,l+1}$  except  $E_{t,l}^{(N_{l,t})}$ . Applying (9), we obtain

$$\begin{aligned} E_{l,l+1}E_{t,l}^{(N_{l,t})}E_{t,l+1}^{(N_{l+1,t})} &= -E_{t,l}^{(N_{l,t}-1)}E_{t,l+1}E_{t,l+1}^{(N_{l+1,t})} + E_{t,l}^{(N_{l,t})}E_{t,l+1}^{(N_{l+1,t})}E_{l,l+1} \\ &= -(N_{l+1,t}+1)E_{t,l}^{(N_{l,t}-1)}E_{t,l+1}^{(N_{l+1,t}+1)} + E_{t,l}^{(N_{l,t})}E_{t,l+1}^{(N_{l+1,t})}E_{l,l+1}. \end{aligned}$$

Hence  $E_{l,l+1}F_t = -(N_{l+1,t}+1)\prod_{1\leq r<t}E_{t,r}^{(N_{r,t}-\delta_{r=l}+\delta_{r=l+1})} + F_tE_{l,l+1}$  and (11) follows.

Now (10) follows from (11) and the equivalence

$$E_{l,l+1}\sum_{N\in\text{UT}(n)}F^{(N)}\mathcal{H}_N \equiv \sum_{N\in\text{UT}(n)}[E_{l,l+1}, F^{(N)}]\mathcal{H}_N \pmod{\mathcal{U}\cdot E_{l,l+1}}.$$

□

We shall identify any graph  $\Gamma$  with vertices  $\mathbf{Z}$  and finitely many edges all having the form  $(s, t)$ , where  $1 \leq s < t \leq n$ , with the matrix of  $\text{UT}(n)$  such that  $N_{s,t}$  equals the number of edges begging at  $s$  and ending at  $t$ .

*Example.* Let  $n = 4$  and  $\Gamma$  have edges  $(1, 2)$ ,  $(1, 4)$ ,  $(2, 3)$ ,  $(3, 4)$ . Then  $\Gamma$  is identified with the matrix

$$\begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

**Definition 25.** Let  $1 \leq a \leq i \leq b < n$ . A sequence of integers satisfies  $\mathcal{O}(a, i, b)$  if this sequence is obtained by inserting the sequences  $a, a+1, \dots, i-1$  and  $b, b-1, \dots, i+1$  into one another, preserving the order in either of them.

**Remark 26.** The only sequence satisfying  $\mathcal{O}(i, i, i)$  is the empty sequence  $\emptyset$ .

*Example.* The sequence  $\underline{1}, 8, \underline{2}, \underline{3}, 7, 6, \underline{4}$  satisfies  $\mathcal{O}(1, 5, 8)$ . We underlined the elements of the sequence  $a, a+1, \dots, i-1$ .

**Lemma 27.** Fix some integers  $1 \leq a_1 < a_2 < \dots < a_q \leq i < b_q < \dots < b_2 < b_1 \leq n$ . Let

$$i_1^{(q)}, \dots, i_{k_q}^{(q)}; \quad i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}; \quad \dots \quad ; \quad i_1^{(1)}, \dots, i_{k_1}^{(1)} \quad (13)$$

be sequences satisfying  $\mathcal{O}(a_q, i, b_q-1)$ ,  $\mathcal{O}(a_{q-1}, i, b_{q-1}-1)$ ,  $\dots$ ,  $\mathcal{O}(a_1, i, b_1-1)$  respectively. For  $\mathcal{H}_N$  as in Lemma 24, we have

$$\begin{aligned} &\llbracket i_1^{(q)}, \dots, i_{k_q}^{(q)}, i, i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}, i, \dots, i_1^{(1)}, \dots, i_{k_1}^{(1)}, i \rrbracket_i \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\ &\sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)} \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+\Gamma}). \end{aligned}$$

*Proof.* We put for brevity

$$\begin{aligned} x &:= \llbracket i_1^{(q)}, \dots, i_{k_q}^{(q)}, i, i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}, i, \dots, i_1^{(1)}, \dots, i_{k_1}^{(1)}, i \rrbracket_i, \\ \mathcal{F} &:= \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q). \end{aligned}$$

We apply induction on the length of the sequence inside the brackets  $\llbracket \rrbracket$  in the formulation of the current lemma (the above formula). If this length

is zero (i.e.  $q = 0$ ), then the required equality holds since  $\llbracket \cdot \rrbracket_i = \theta_{\omega_i}$ ,  $\mathcal{F}_i(\cdot)$  consists of the empty flow  $\Gamma_\emptyset$ , which is identified with the zero matrix, and  $\text{sgn}_i(\Gamma_\emptyset) = 1$ .

Now suppose that the equality in the formulation of the current lemma holds. Take any  $l = 1, \dots, n-1$  and set

$$y := [\eta_{\alpha_l}, x] = \left\llbracket l, i_1^{(q)}, \dots, i_{k_q}^{(q)}, i, i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}, i, \dots, i_1^{(1)}, \dots, i_{k_1}^{(1)}, i \right\rrbracket_i. \quad (14)$$

Applying (10) and the inductive hypothesis, we get

$$\begin{aligned} & \eta_{\alpha_l} x \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\ & \eta_{\alpha_l} \left( \sum_{N \in \text{UT}(n)} F^{(N)} \left[ \sum_{\Gamma \in \mathcal{F}} \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+\Gamma}) \right] \right) = \\ & \sum_{N \in \text{UT}(n)} F^{(N)} \left( \sum_{1 \leq s < l} N_{s,l} \left[ \sum_{\Gamma \in \mathcal{F}} \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+E_{s,l+1}-E_{s,l}+\Gamma}) \right] \right. \\ & \quad \left. + (H_l - N^l + N^{l+1}) \left[ \sum_{\Gamma \in \mathcal{F}} \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+E_{l,l+1}+\Gamma}) \right] \right. \\ & \quad \left. - \sum_{l+1 < t \leq n} N_{l+1,t} \left[ \sum_{\Gamma \in \mathcal{F}} \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+E_{l,t}-E_{l+1,t}+\Gamma}) \right] \right) = \\ & \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}} \left[ \sum_{1 \leq s < l} \text{sgn}_i(\Gamma) N_{s,l} \theta_{\omega_i}(\mathcal{H}_{N+E_{s,l+1}-E_{s,l}+\Gamma}) \right. \\ & \quad \left. + \text{sgn}_i(\Gamma) (H_l - N^l + N^{l+1}) \theta_{\omega_i}(\mathcal{H}_{N+E_{l,l+1}+\Gamma}) \right. \\ & \quad \left. - \sum_{l+1 < t \leq n} \text{sgn}_i(\Gamma) N_{l+1,t} \theta_{\omega_i}(\mathcal{H}_{N+E_{l,t}-E_{l+1,t}+\Gamma}) \right]. \end{aligned}$$

On the other hand, we get

$$\begin{aligned} & x \eta_{\alpha_l} \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\ & x \left( \sum_{N \in \text{UT}(n)} F^{(N)} \left[ \sum_{1 \leq s < l} N_{s,l} \mathcal{H}_{N+E_{s,l+1}-E_{s,l}} \right. \right. \\ & \quad \left. \left. + (H_l - N^l + N^{l+1}) \mathcal{H}_{N+E_{l,l+1}} - \sum_{l+1 < t \leq n} N_{l+1,t} \mathcal{H}_{N+E_{l,t}-E_{l+1,t}} \right] \right) = \\ & \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}} \left[ \sum_{1 \leq s < l} \text{sgn}_i(\Gamma) (N+\Gamma)_{s,l} \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{s,l+1}-E_{s,l}}) \right. \\ & \quad \left. + \text{sgn}_i(\Gamma) (\theta_{\omega_i}(H_l) - (N+\Gamma)^l + (N+\Gamma)^{l+1}) \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{l,l+1}}) \right. \\ & \quad \left. - \sum_{l+1 < t \leq n} \text{sgn}_i(\Gamma) (N+\Gamma)_{l+1,t} \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{l,t}-E_{l+1,t}}) \right]. \end{aligned}$$

Subtracting the latter expression from the former, we obtain (recall the definition of  $\theta_\delta$  in Section 2.4)

$$\begin{aligned}
& y \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\
& \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}} \left[ \sum_{1 \leq s < l} -\text{sgn}_i(\Gamma) \Gamma_{s,l} \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{s,l+1}-E_{s,l}}) \right. \\
& \quad \left. + \text{sgn}_i(\Gamma) (-\delta_{i=l} + \Gamma^l - \Gamma^{l+1}) \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{l,l+1}}) + \right. \\
& \quad \left. \sum_{l+1 < t \leq n} \text{sgn}_i(\Gamma) \Gamma_{l+1,t} \theta_{\omega_i}(\mathcal{H}_{N+\Gamma+E_{l,t}-E_{l+1,t}}) \right].
\end{aligned}$$

We denote by  $S_1(\Gamma)$ ,  $S_2(\Gamma)$  and  $S_3(\Gamma)$  the first, the second and the third summands respectively in the square brackets in the above formula.

Now we want to choose  $l$  so that the sequence in the brackets  $\llbracket \cdot \rrbracket$  of (14) could appear in the brackets  $\llbracket \cdot \rrbracket$  in the formulation of the current lemma for different sequences (13) and possibly different  $q$  (Case 2).

*Case 1:*  $q \geq 1$  and  $1 \leq l = a_q - 1 \notin \{a_1, \dots, a_{q-1}\}$ . The values of  $S_1(\Gamma)$ ,  $S_2(\Gamma)$  and  $S_3(\Gamma)$  are given in the table below.

|               | $a_q - 1$ is no transit point of $\Gamma$                              | $a_q - 1$ is a transit point of $\Gamma$                               |
|---------------|------------------------------------------------------------------------|------------------------------------------------------------------------|
| $S_1(\Gamma)$ | 0                                                                      | $-\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_1(\Gamma)})$ |
| $S_2(\Gamma)$ | $-\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_2(\Gamma)})$ | 0                                                                      |
| $S_3(\Gamma)$ | $\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_3(\Gamma)})$  | $\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_3(\Gamma)})$  |

Table 4

In each row of this table (except the top row), the element of  $\mathcal{U}$  in the left column equals either the element in the middle column or the element in the right column depending on which condition in the top row of this table is satisfied for  $\Gamma$ .

To verify this table, one should note that in the present case  $l < i$ ,  $\Gamma^{l+1} = 1$  (since  $l+1 = a_q$  is a source of  $\Gamma$ ) and  $\Gamma^l = 0$  if  $a_q - 1$  is no transit point of  $\Gamma$  and  $\Gamma^l = 1$  otherwise. By Lemmas 12–14, every nonzero cell of this table (not in the top row and not in the left column) that is not the bottom rightmost yields a flow of  $\mathcal{F}_i(a_1, \dots, a_{q-1}, a_q - 1; b_1, \dots, b_q)$  as the second summand in the subscript of  $\mathcal{H}$ . Therefore, applying Lemmas 12–14 and Corollary 19, we obtain

$$\begin{aligned}
& y \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\
& \sum_{N \in \text{UT}(n)} F^{(N)} \left( \sum \left\{ \text{sgn}_i(L_k(\Gamma)) \theta_{\omega_i}(\mathcal{H}_{N+L_k(\Gamma)}) \mid \Gamma \in \mathcal{F} \ \& \ k = 1, 2, 3 \right. \right. \\
& \quad \left. \& \ L_k(\Gamma) \text{ is well-defined and a flow} \right\} + \\
& \quad \left. \sum \left\{ \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_3(\Gamma)}) \mid \Gamma \in \mathcal{F} \ \& \ a_q - 1 \text{ is a transit point of } \Gamma \right\} \right) =
\end{aligned}$$

$$\sum_{N \in \text{UT}(n)} F^{(N)} \left( \sum_{\Gamma' \in \mathcal{F}_i(a_1, \dots, a_{q-1}, a_q-1; b_1, \dots, b_q)} \text{sgn}_i(\Gamma') \theta_{\omega_i}(\mathcal{H}_{N+\Gamma'}) + \right. \\ \left. \sum \left\{ \text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+L_3(\Gamma)}) \mid \Gamma \in \mathcal{F} \text{ \& } a_q-1 \text{ is a transit point of } \Gamma \right\} \right).$$

By Lemma 14, the last summand of the above formula equals zero. Thus we have proved the current lemma for the sequences

$$a_q-1, i_1^{(q)}, \dots, i_{k_q}^{(q)}; \quad i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}; \quad \dots \quad ; \quad i_1^{(1)}, \dots, i_{k_1}^{(1)} \quad (15)$$

satisfying conditions  $\mathcal{O}(a_q-1, i, b_q-1)$ ,  $\mathcal{O}(a_{q-1}, i, b_{q-1}-1)$ ,  $\dots$ ,  $\mathcal{O}(a_1, i, b_1-1)$  respectively.

*Case 2:*  $a_q < i$ ,  $i+1 < b_q$  if  $q \geq 1$  and  $l = i$ . The values of  $S_1(\Gamma)$ ,  $S_2(\Gamma)$  and  $S_3(\Gamma)$  are given in the table below.

|               | $i$ is no transit point of $\Gamma$                                    | $i$ is a transit point of $\Gamma$                                     |
|---------------|------------------------------------------------------------------------|------------------------------------------------------------------------|
| $S_1(\Gamma)$ | 0                                                                      | $-\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+M_1(\Gamma)})$ |
| $S_2(\Gamma)$ | $-\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+M_2(\Gamma)})$ | 0                                                                      |
| $S_3(\Gamma)$ | 0                                                                      | 0                                                                      |

Table 5

This table should be interpreted in the same way as Table 4.

To verify this table, one should note that in the present case  $\Gamma_{l+1,t} = 0$  for any  $t \in \mathbf{Z}$ ,  $\Gamma^{l+1} = 0$ ,  $\Gamma^l = 0$  if  $i$  is no transit point of  $\Gamma$  and  $\Gamma^l = 1$  otherwise. Therefore, applying Lemmas 15 and 16 and Corollary 21, we obtain

$$y \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\ \sum_{N \in \text{UT}(n)} F^{(N)} \sum \left\{ \text{sgn}_i(M_k(\Gamma)) \theta_{\omega_i}(\mathcal{H}_{N+M_k(\Gamma)}) \mid \Gamma \in \mathcal{F} \text{ \& } k = 1, 2 \right. \\ \left. \text{\& } M_k(\Gamma) \text{ is well-defined} \right\} = \\ \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q, i; b_1, \dots, b_q, i+1)} \text{sgn}_i(\Gamma') \theta_{\omega_i}(\mathcal{H}_{N+\Gamma'}).$$

Thus we have proved the current lemma for the  $q+1$  sequences

$$\emptyset; \quad i_1^{(q)}, \dots, i_{k_q}^{(q)}; \quad i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}; \quad \dots \quad ; \quad i_1^{(1)}, \dots, i_{k_1}^{(1)} \quad (16)$$

satisfying conditions  $\mathcal{O}(i, i, i)$ ,  $\mathcal{O}(a_q, i, b_q-1)$ ,  $\mathcal{O}(a_{q-1}, i, b_{q-1}-1)$ ,  $\dots$ ,  $\mathcal{O}(a_1, i, b_1-1)$  respectively (see Remark 26).

*Case 3:*  $q \geq 1$ ,  $b_q+1 \notin \{b_{q-1}, \dots, b_1\}$  and  $l = b_q$ . We have  $i < l$ ,  $\Gamma_{l+1,t} = 0$  for any  $t \in \mathbf{Z}$ ,  $\Gamma^{l+1} = 0$ ,  $\Gamma^l = 0$ . Hence  $S_1(\Gamma) = -\text{sgn}_i(\Gamma) \theta_{\omega_i}(\mathcal{H}_{N+R(\Gamma)})$ ,  $S_2(\Gamma) = 0$  and  $S_3(\Gamma) = 0$ . Therefore, applying Lemma 17 and Corollary 23, we obtain

$$y \left( \sum_{N \in \text{UT}(n)} F^{(N)} \mathcal{H}_N \right) = \\ \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}} \text{sgn}_i(R(\Gamma)) \theta_{\omega_i}(\mathcal{H}_{N+R(\Gamma)}) = \\ \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_{q-1}, b_q+1)} \text{sgn}_i(\Gamma') \theta_{\omega_i}(\mathcal{H}_{N+\Gamma'}).$$

Thus we have proved the current lemma for the sequences

$$b_q, i_1^{(q)}, \dots, i_{k_q}^{(q)}; \quad i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}; \quad \dots \quad ; \quad i_1^{(1)}, \dots, i_{k_1}^{(1)} \quad (17)$$

satisfying conditions  $\mathcal{O}(a_q, i, b_q)$ ,  $\mathcal{O}(a_{q-1}, i, b_{q-1} - 1)$ ,  $\dots$ ,  $\mathcal{O}(a_1, i, b_1 - 1)$  respectively.

To finish the proof, it suffices to notice that any sequence of sequences in the hypothesis of the current lemma can be obtained by passing from (13) to (15) or (16) or (17).  $\square$

**Definition 28.** It follows from Lemma 27 that the operator  $\llbracket i_1^{(q)}, \dots, i_{k_q}^{(q)}, i, i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}, i, \dots, i_1^{(1)}, \dots, i_{k_1}^{(1)}, i \rrbracket_i$  depends only on the integers  $a_1, \dots, a_q, b_1, \dots, b_q$  and  $i$ . We, therefore, denote it by  $\xi_i(a_1, \dots, a_q; b_1, \dots, b_q)$ .

**3.4. Tableaux.** A *composition* of length  $n$  is a sequence  $\lambda = (\lambda_1, \dots, \lambda_n)$  such that  $\lambda_1, \dots, \lambda_n \in \mathbf{Z}^+$ . If, moreover,  $\lambda_1 \geq \dots \geq \lambda_n$  then  $\lambda$  is called a *partition*. The *diagram* of a composition  $\lambda$  is the set

$$[\lambda] = \{(i, j) \in \mathbf{Z}^2 \mid 1 \leq i \leq n \text{ and } 1 \leq j \leq \lambda_i\}.$$

We shall think of  $[\lambda]$  as an array of boxes. For example, if  $\lambda = (5, 3, 2, 0)$  then

$$[\lambda] = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array}$$

A  $\lambda$ -*tableau*, where  $\lambda$  is a composition, is a function  $T : [\lambda] \rightarrow \{1, \dots, n\}$ , which we regard as the diagram  $[\lambda]$  filled with integers in  $\{1, \dots, n\}$ . A  $\lambda$ -tableau  $T$  is called *row standard* if its entries weakly increase along the rows, that is  $T(i, j) \leq T(i, j')$  if  $j < j'$ . A  $\lambda$ -tableau  $T$  is called *regular row standard* if it is row standard and every entry in row  $i$  of  $T$  is at least  $i$ . Finally, if  $\lambda$  is partition, then a  $\lambda$ -tableau  $T$  is called *standard* if it is row standard and its entries strictly increase down the columns, that is  $T(i, j) < T(i', j)$  if  $i < i'$ . For example,

$$T = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 2 & 3 & 4 \\ \hline 2 & 3 & 3 & & \\ \hline 3 & 4 & & & \\ \hline \end{array}$$

is a standard  $(5, 3, 2, 0)$ -tableau. Here  $n = 4$ .

For any regular row standard  $\lambda$ -tableau  $T$ , we define  $\text{Mat}_T$  to be the matrix of  $\text{UT}(n)$  whose  $(i, j)$ -entry, where  $i < j$ , is the number of entries  $j$  in row  $i$  of  $T$  and set  $F_T := F^{(\text{Mat}_T)}$ . For  $n = 4$  and  $T$  as in the above example, we have

$$\text{Mat}_T = \begin{pmatrix} 0 & 2 & 1 & 1 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad F_T = E_{2,1}^{(2)} E_{3,1} E_{3,2}^{(2)} E_{4,1} E_{4,3}.$$

We say that a composition  $\lambda = (\lambda_1, \dots, \lambda_n)$  is *coherent* with a weight  $c_1\omega_1 + \dots + c_{n-1}\omega_{n-1}$  if  $c_i = \lambda_i - \lambda_{i+1}$  for any  $i = 1, \dots, n-1$ .

**Proposition 29** ([5]). *Let  $\omega$  be a dominant weight and  $\lambda$  be any partition coherent with  $\omega$ . Then the vectors  $F_T e_\omega^+$ , where  $T$  is a standard  $\lambda$ -tableau, form a basis of  $\Delta(\omega)$ .*

To deal with compositions, it is convenient to introduce the following sequences of length  $n$ :  $\varepsilon_i := (0, \dots, 0, 1, 0, \dots, 0)$  with 1 at position  $i$  and  $\alpha(i, j) = \varepsilon_i - \varepsilon_j$ . Here and in what follows, the sequences of  $\mathbf{Z}^n$  are added and subtracted componentwise.

For the rest of Sections 3.4 – 3.6, we fix integers  $a_1, \dots, a_q, b_1, \dots, b_q, i$  such that

$$1 \leq a_1 < a_2 < \dots < a_q \leq i < b_q < \dots < b_2 < b_1 \leq n. \quad (18)$$

Let  $T$  be a regular row standard  $\lambda$ -tableau and  $\Gamma$  be a flow of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ . We want to construct the regular row standard tableau  $\sigma_{\Gamma, i}(T)$  so that

( $\sigma$ -1)  $\text{Mat}_T - \Gamma = \text{Mat}_{\sigma_{\Gamma, i}(T)}$ ;

( $\sigma$ -2)  $\sigma_{\Gamma, i}(T)$  is a  $(\lambda_1, \dots, \lambda_i, \lambda_{i+1} + 1, \dots, \lambda_n + 1)$ -tableau.

We obtain  $\sigma_{\Gamma, i}(T)$  from  $T$  as follows

- (i) for every edge  $(s, t)$  of  $\Gamma$ , we replace one  $t$  with  $s$  in row  $s$ ;
- (ii) to every row  $k = i + 1, \dots, n$ , we add one  $k$ ;
- (iii) in the resulting tableau, we order the elements in rows to obtain a row standard tableau (automatically regular).

One can see that (i) ensures ( $\sigma$ -1) and (ii) ensures ( $\sigma$ -2), while (iii) makes the tableau row standard and eliminates the uncertainty stemming from the choice of the elements being removed and added in steps (i) and (ii).

Obviously, it is not always possible to executed all replacements in step (i), in which case  $\sigma_{\Gamma, i}(T)$  is not well-defined. This happens if and only if for some edge  $(s, t)$  of  $\Gamma$ , row  $s$  of  $T$  does not contain  $t$  as an entry, i.e. if and only if  $\Gamma_{s, t} = 1$  and  $(\text{Mat}_T)_{s, t} = 0$  for some  $s$  and  $t$ . Thus we have proved the following property:

( $\sigma$ -3)  $\sigma_{\Gamma, i}(T)$  is well-defined if and only if all entries of  $\text{Mat}_T - \Gamma$  are nonnegative.

For example, take  $n = 4$ ,  $i = 2$ ,  $\Gamma \in \mathcal{F}_2(1, 2; 4, 3)$  with edges  $(1, 4)$ ,  $(2, 3)$  and  $T$  as in the above example. Then

$$\begin{array}{ccc}
 T = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 2 & 3 & 3 & \\ \hline 3 & 4 & & \\ \hline \end{array} & \xrightarrow{\text{step (i)}} & \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 2 & 2 & 3 & \\ \hline 3 & 4 & & \\ \hline \end{array} & \xrightarrow{\text{step (ii)}} & \\
 \\
 \begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 3 \\ \hline 2 & 2 & 3 & \\ \hline 3 & 4 & 3 & \\ \hline 4 & & & \\ \hline \end{array} & \xrightarrow{\text{step (iii)}} & \begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 2 \\ \hline 2 & 2 & 3 & \\ \hline 3 & 3 & 4 & \\ \hline 4 & & & \\ \hline \end{array} & = & \sigma_{\Gamma, i}(T)
 \end{array}$$

On the other hand, if we take  $\Gamma' \in \mathcal{F}_2(2; 4)$  with edge  $(2, 4)$ , then  $\sigma_{\Gamma', i}(T)$  is not well-defined, since there is no 4 in row 2 of  $T$  and step (i) can not be executed. We also have  $(\text{Mat}_T - \Gamma')_{2, 4} = -1$  in accordance with ( $\sigma$ -3).

**Corollary 30.** *Let  $T$  be a regular row standard  $\lambda$ -tableau. Then  $\xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F_T)$  is the sum of  $\text{sgn}_i(\Gamma)F_{\sigma_{\Gamma,i}(T)}$  over all flows  $\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $\sigma_{\Gamma,i}(T)$  is well-defined.*

*Proof.* By Lemma 27, we have (recall Definition 28)

$$\begin{aligned} \xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F_T) &= \\ \xi_i(a_1, \dots, a_q; b_1, \dots, b_q) \left( \sum_{N \in \text{UT}(n)} F^{(N)} \delta_{N=\text{Mat}_T} \right) &= \\ \sum_{N \in \text{UT}(n)} F^{(N)} \sum_{\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)} \text{sgn}_i(\Gamma) \delta_{N+\Gamma=\text{Mat}_T} &= \\ \sum_{\Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)} \text{sgn}_i(\Gamma) F^{(\text{Mat}_T - \Gamma)}. \end{aligned}$$

Clearly,  $F^{(\text{Mat}_T - \Gamma)} = 0$  if at least one entry of  $\text{Mat}_T - \Gamma$  is negative. Therefore, by  $(\sigma-1)$  and  $(\sigma-3)$ , we obtain

$$\begin{aligned} \xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F_T) &= \\ \sum \left\{ \text{sgn}_i(\Gamma) F^{(\text{Mat}_T - \Gamma)} \mid \Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q) \text{ \& } \right. & \\ \left. \text{all entries of } \text{Mat}_T - \Gamma \text{ are nonnegative} \right\} &= \\ \sum \left\{ \text{sgn}_i(\Gamma) F_{\sigma_{\Gamma,i}(T)} \mid \Gamma \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q) \text{ \& } \sigma_{\Gamma,i}(T) \text{ is well-defined} \right\}. \end{aligned}$$

□

**3.5. Comparison of tableaux.** We make the following convention on orders. The symbol  $\leq$  (symbol  $<$ ) will always denote a nonstrict (resp. strict) partial order on a set  $S$ , i.e. a transitive binary relation of  $S$  such that  $x \leq x$  and  $x \leq y \text{ \& } y \leq x \Rightarrow x = y$  for any  $x, y \in S$  (resp.  $x < y \text{ \& } y < x$  for no  $x, y \in S$ ). We also assume that for any  $x, y \in S$ , there holds

$$\begin{aligned} x \leq y &\Leftrightarrow x < y \text{ or } x = y, & x \leq y &\Leftrightarrow y \geq x, \\ x < y &\Leftrightarrow x \leq y \text{ \& } x \neq y, & x < y &\Leftrightarrow y > x. \end{aligned}$$

Consider two sequences  $x = (x_1, \dots, x_m)$  and  $y = (y_1, \dots, y_m)$  of elements of  $S$ . We say that  $x$  is smaller than  $y$  in the antilexicographic order if there exists  $i = 1, \dots, m$  such that  $x_i < y_i$  and  $x_j = y_j$  for  $i < j \leq m$ .

The set of all compositions of length  $n$  is endowed with the following partial order called *dominance order*. For two compositions  $\lambda = (\lambda_1, \dots, \lambda_n)$  and  $\mu = (\mu_1, \dots, \mu_n)$ , we write  $\lambda \leq \mu$  if  $\sum_{j=1}^i \lambda_j \leq \sum_{j=1}^i \mu_j$  for any  $i = 1, \dots, n$ .

Let  $\mu$  be a composition of length  $n$  and  $Q$  be a  $\mu$ -tableau. Then we set  $\text{shape}(Q) := \mu$ . Given a composition  $\lambda$  of length  $n$ , a row standard  $\lambda$ -tableau  $T$  and an integer  $m = 1, \dots, n$ , we denote by  $T[m]$  the tableau obtained from  $T$  by removing all nodes with entry greater than  $m$ . Clearly,  $T[m]$  is a  $\mu$ -tableau for some composition  $\mu$  of length  $n$ . Therefore, we define

$$\text{chain}(T) := (\text{shape}(T[1]), \text{shape}(T[2]), \dots, \text{shape}(T[n])).$$

For another row standard  $\lambda$ -tableau  $S$ , we write  $T < S$  if  $\text{chain}(T)$  is smaller than  $\text{chain}(S)$  in the antilexicographic order with dominance order on components. We also refer the reader to [1, Section 2.2] for a description of this order.

It is convenient to define the sum of row standard tableaux  $S$  and  $T$  as the row standard tableau  $S + T$ , whose  $i$ th row is the result of gluing and reordering the  $i$ th rows of  $S$  and  $T$ . Obviously, we have

$$(S + T)[m] = S[m] + T[m], \quad \text{shape}(S + T) = \text{shape}(S) + \text{shape}(T). \quad (19)$$

*Example.* Let  $n = 5$ . Then we have

$$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 4 & 5 \\ \hline 2 & 2 & 4 & \\ \hline 5 & & & \\ \hline 5 & & & \\ \hline \end{array} + \begin{array}{|c|c|} \hline 3 & \\ \hline 3 & 5 \\ \hline 4 & 5 \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5 \\ \hline 2 & 2 & 3 & 4 & 5 \\ \hline 4 & 5 & 5 & & \\ \hline 5 & & & & \\ \hline \end{array}$$

Applying shape to each term gives (cf. (19))

$$(4, 3, 1, 1, 0) + (1, 2, 2, 0, 0) = (5, 5, 3, 1, 0).$$

To understand how  $\text{chain}(T)$  changes when  $T$  is replaced with  $\sigma_{\Gamma,i}(T)$ , we introduce the following notation. For a flow  $\Gamma$  and an integer  $m$ , we set

$$\nu_m(\Gamma) := \sum \{\varepsilon_s \mid (s, t) \text{ is an edge of } \Gamma \text{ and } s \leq m < t\}.$$

**Lemma 31.** *Let  $\Gamma$  be a flow of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and  $m = 1, \dots, n$ . Then we have*

$$\text{shape}(\sigma_{\Gamma,i}(T)[m]) = \text{shape}(T[m]) + \nu_m(\Gamma) + \sum_{k=i+1}^m \varepsilon_k$$

if  $\sigma_{\Gamma,i}$  is applicable to  $T$ .

*Proof.* Let  $S$  be the tableau obtained from  $T$  in the same way as  $\sigma_{\Gamma,i}(T)$  but without step (ii) and  $Q$  be the  $(0^i, 1^{n-i})$ -tableau having one entry  $k$  in each row  $k = i + 1, \dots, n$ . We clearly have  $\sigma_{\Gamma,i}(T) = S + Q$ . It follows from (19) that  $\text{shape}(\sigma_{\Gamma,i}(T)[m]) = \text{shape}(S[m]) + \text{shape}(Q[m])$ . It is easy to see that  $\text{shape}(Q[m]) = \sum_{k=i+1}^m \varepsilon_k$ .

Let us divide each row of  $T$  and  $S$  into the left part containing entries  $\leq m$  and the right part containing entries  $> m$ . Thus the right parts are deleted and the left parts remain under  $T \mapsto T[m]$  and  $S \mapsto S[m]$ .

Obviously, the left part of row  $s$  of  $S$  has different length than the left part of row  $s$  of  $T$  if and only if  $s \leq m < t$  for some edge  $(s, t)$  of  $\Gamma$  (i.e., in step (i), an entry of the right part of row  $s$  of  $T$  is replaced by an entry that will go to the left part of row  $s$  of  $S$ ). In that case, the left part of row  $s$  of  $S$  is longer by one than left part of row  $s$  of  $T$ , which proves  $\text{shape}(S[m]) = \text{shape}(T[m]) + \nu_m(\Gamma)$  and the lemma.  $\square$

**Proposition 32** (Straightening rule [1, Theorem A.4]). *If  $\lambda$  is a partition coherent with a dominant weight  $\omega$  and  $T$  is a regular row standard but not standard  $\lambda$ -tableau, then  $F_{Te_\omega^+}$  can be rewritten as a linear combination of vectors  $F_{Se_\omega^+}$ , where  $S$  is a standard  $\lambda$ -tableau such that  $S > T$ .*



**3.6. Comparison of flows.** We are going to introduce a partial order on  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and show its relation to the order of Section 3.5. We start with an obvious observation.

**Proposition 33.** *Let  $\Gamma$  be a nonempty flow with sources  $\mathbf{a}_1, \dots, \mathbf{a}_q$  and sinks  $\mathbf{b}_1, \dots, \mathbf{b}_q$ . Take the edge  $(\hat{s}, \hat{t})$  of  $\Gamma$  such that  $\hat{t} > t$  for any other edge  $(s, t)$  of  $\Gamma$ . We have  $\hat{t} = \mathbf{b}_j$  for some  $j = 1, \dots, q$ . The graph  $\tilde{\Gamma}$  obtained from  $\Gamma$  by removing  $(\hat{s}, \hat{t})$  is again a flow with*

- (i) *sources  $\mathbf{a}_1, \dots, \mathbf{a}_q$  and sinks  $\mathbf{b}_1, \dots, \mathbf{b}_{j-1}, \hat{s}, \mathbf{b}_{j+1}, \dots, \mathbf{b}_q$  if  $\hat{s} \notin \{\mathbf{a}_1, \dots, \mathbf{a}_q\}$ ;*
- (ii) *sources  $\mathbf{a}_1, \dots, \mathbf{a}_{k-1}, \mathbf{a}_{k+1}, \dots, \mathbf{a}_q$  and sinks  $\mathbf{b}_1, \dots, \mathbf{b}_{j-1}, \mathbf{b}_{j+1}, \dots, \mathbf{b}_q$  if  $\hat{s} = \mathbf{a}_k$ .*

Note that in this proposition the numbers  $\mathbf{a}_1, \dots, \mathbf{a}_q, \mathbf{b}_1, \dots, \mathbf{b}_q$  do not have to satisfy (18) unlike  $a_1, \dots, a_q, b_1, \dots, b_q$ .

Let  $\Gamma$  and  $\Gamma'$  be distinct flows of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  and

$$(s_1, t_1), \dots, (s_r, t_r); \quad (s'_1, t'_1), \dots, (s'_{r'}, t'_{r'}) \quad (20)$$

be the edges of  $\Gamma$  and  $\Gamma'$  respectively ordered so that  $t_1 > \dots > t_r$  and  $t'_1 > \dots > t'_{r'}$ .

Proposition 33 explains why one sequence of (20) can not be a proper beginning of the other. Indeed, assume that  $r < r'$  and  $(s_k, t_k) = (s'_k, t'_k)$  for all  $k = 1, \dots, r$ . Let  $\tilde{\Gamma}$  and  $\tilde{\Gamma}'$  be the graphs obtained from  $\Gamma$  and  $\Gamma'$  respectively by removing the edges  $(s_1, t_1), \dots, (s_r, t_r)$ . Then by Proposition 33 applied  $r$  times,  $\tilde{\Gamma}$  and  $\tilde{\Gamma}'$  are flows with the same sources and sinks. To understand this, note that in (i) and (ii) of Proposition 33, new sources and sinks depend only on  $a_1, \dots, a_q, b_1, \dots, b_q$  and the edge  $(\hat{s}, \hat{t})$  being removed. However  $\tilde{\Gamma}$  is the empty flow, while  $\tilde{\Gamma}'$  is not, since  $r' > r$ . This is a contradiction.

By what we have just proved, there exists an index  $l = 1, \dots, \min\{r, r'\}$  such that  $(s_k, t_k) = (s'_k, t'_k)$  for  $k < l$  and  $(s_l, t_l) \neq (s'_l, t'_l)$ . Let  $\Gamma_0$  and  $\Gamma'_0$  be the graphs obtained from  $\Gamma$  and  $\Gamma'$  respectively by removing the edges  $(s_1, t_1), \dots, (s_{l-1}, t_{l-1})$ . By Proposition 33 applied  $l-1$  times,  $\Gamma_0$  and  $\Gamma'_0$  are flows with the same sources and sinks. Hence  $t_l = t'_l$ , since  $t_l$  is the maximal sink of  $\Gamma_0$  and  $t'_l$  is the maximal sink of  $\Gamma'_0$ . We write  $\Gamma < \Gamma'$  if  $s_l > s'_l$ .

We still need to prove that the relation on  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  we have defined is a linear order. Clearly, there exists a linear order  $<$  on  $\mathbf{Z}^2$  such that

$$(s, t) < (s', t) \text{ if and only if } s > s'.$$

Then it is obvious that  $\Gamma < \Gamma'$  if and only if  $(s_1, t_1), \dots, (s_r, t_r)$  is lexicographically smaller than  $(s'_1, t'_1), \dots, (s'_{r'}, t'_{r'})$  with respect to the chosen order on  $\mathbf{Z}^2$ .

*Example.* Consider the flows  $\Gamma$  and  $\Gamma'$  of  $\mathcal{F}_4(1, 3; 6, 5)$  with edges

$$(3, 6), (4, 5), (2, 4), (1, 2); \quad (3, 6), (4, 5), (1, 4)$$

respectively. The first two edges are the same in both graphs. Therefore looking at the third edges  $(2, 4)$  and  $(1, 4)$ , we conclude  $\Gamma < \Gamma'$ .

**Lemma 34.** *Let  $\Gamma$  and  $\Gamma'$  be flows of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $\Gamma < \Gamma'$ . Then  $(\nu_1(\Gamma), \dots, \nu_n(\Gamma))$  is smaller than  $(\nu_1(\Gamma'), \dots, \nu_n(\Gamma'))$  in the antilexicographic order with the dominance order on components.*

*Proof.* We use the notation introduced before this lemma. Recall that we have  $(s_k, t_k) = (s'_k, t'_k)$  for  $k < l$ ,  $t_l = t'_l$  and  $s_l > s'_l$ . Take any integer  $m$  such that  $t_l - 1 \leq m \leq n$ . We denote by  $N_m$  and  $N'_m$  the numbers of elements  $> m$  in the sequences  $t_1, \dots, t_r$  and  $t'_1, \dots, t'_{r'}$  respectively. Since  $t_{l+1} \leq t_l - 1 \leq m$  if  $l < r$  and  $t'_{l+1} \leq t'_l - 1 \leq m$  if  $l < r'$ , we obtain that  $N_m$  and  $N'_m$  are the numbers of elements  $> m$  in the sequences  $t_1, \dots, t_l$  and  $t'_1, \dots, t'_l$  respectively. However these sequences are equal. Thus  $N_m = N'_m$ .

If  $m \geq t_l$  then  $N_m < l$  and  $\nu_m(\Gamma) = \nu_m(\Gamma') = \sum_{k=1}^{N_m} \delta_{s_k \leq m} \varepsilon_{s_k}$ . On the other hand,  $N_{t_l-1} = l$  and

$$\begin{aligned} \nu_{t_l-1}(\Gamma') - \nu_{t_l-1}(\Gamma) &= \sum_{k=1}^l \delta_{s'_k \leq t_l-1} \varepsilon_{s'_k} - \sum_{k=1}^l \delta_{s_k \leq t_l-1} \varepsilon_{s_k} = \\ \varepsilon_{s'_l} - \varepsilon_{s_l} &= \alpha(s'_l, s_l) > 0. \end{aligned}$$

To check the last inequality, see the definition of the dominance order in Section 3.5.  $\square$

For  $\Gamma, \Gamma' \in \mathcal{F}_4(1, 3; 6, 5)$  as in the above example, we have

$$\begin{aligned} \nu_6(\Gamma) &= 0, & \nu_5(\Gamma) &= \varepsilon_3, & \nu_4(\Gamma) &= \varepsilon_3 + \varepsilon_4, & \nu_3(\Gamma) &= \varepsilon_2 + \varepsilon_3, \\ \nu_6(\Gamma') &= 0, & \nu_5(\Gamma') &= \varepsilon_3, & \nu_4(\Gamma') &= \varepsilon_3 + \varepsilon_4, & \nu_3(\Gamma') &= \varepsilon_1 + \varepsilon_3, \end{aligned}$$

and  $\nu_3(\Gamma) < \nu_3(\Gamma')$  as predicted by Lemma 34.

**Lemma 35.** *Let  $T$  and  $S$  be regular row standard tableaux,  $\Gamma$  and  $\Gamma'$  be flows of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  such that  $\Gamma \leq \Gamma'$  and  $\sigma_{\Gamma', i}(S) \leq \sigma_{\Gamma, i}(T)$ . Then we have  $S \leq T$  and, moreover,  $S < T$  if  $\Gamma < \Gamma'$ .*

*Proof.* By Lemma 31, for any  $m = 1, \dots, n$ , we have

$$\text{shape}(\sigma_{\Gamma, i}(T)[m]) - \text{shape}(\sigma_{\Gamma', i}(S)[m]) = \text{shape}(T[m]) - \text{shape}(S[m])$$

if  $\Gamma = \Gamma'$ . Therefore,  $\sigma_{\Gamma', i}(S) \leq \sigma_{\Gamma, i}(T)$  immediately implies  $S \leq T$ .

Therefore, we consider only the case  $\Gamma < \Gamma'$ . By Lemma 34, the sequence  $(\nu_1(\Gamma), \dots, \nu_n(\Gamma))$  is smaller than the sequence  $(\nu_1(\Gamma'), \dots, \nu_n(\Gamma'))$  in the antilexicographic order. This means that there is some  $x = 1, \dots, n$  such that  $\nu_x(\Gamma) < \nu_x(\Gamma')$  and  $\nu_m(\Gamma) = \nu_m(\Gamma')$  for  $x < m \leq n$ .

Let  $y = 1, \dots, n$  be the number such that  $\text{shape}(\sigma_{\Gamma', i}(S)[y]) < \text{shape}(\sigma_{\Gamma, i}(T)[y])$  and  $\text{shape}(\sigma_{\Gamma', i}(S)[m]) = \text{shape}(\sigma_{\Gamma, i}(T)[m])$  for  $y < m \leq n$  if  $\sigma_{\Gamma', i}(S) < \sigma_{\Gamma, i}(T)$  and let  $y := -\infty$  if  $\sigma_{\Gamma', i}(S) = \sigma_{\Gamma, i}(T)$ .

*Case  $y > x$ .* For  $m > y$ , we have  $m > x$  and thus  $\nu_m(\Gamma) = \nu_m(\Gamma')$ . Therefore, by Lemma 31, we get

$$0 = \text{shape}(\sigma_{\Gamma, i}(T)[m]) - \text{shape}(\sigma_{\Gamma', i}(S)[m]) = \text{shape}(T[m]) - \text{shape}(S[m]).$$

On the other hand,  $\nu_y(\Gamma) = \nu_y(\Gamma')$  and Lemma 31 yields

$$0 < \text{shape}(\sigma_{\Gamma, i}(T)[y]) - \text{shape}(\sigma_{\Gamma', i}(S)[y]) = \text{shape}(T[y]) - \text{shape}(S[y]).$$

*Case  $y \leq x$ .* For  $m > x$ , we have  $\nu_m(\Gamma) = \nu_m(\Gamma')$  and  $m > y$ . Therefore, by Lemma 31, we get

$$0 = \text{shape}(\sigma_{\Gamma, i}(T)[m]) - \text{shape}(\sigma_{\Gamma', i}(S)[m]) = \text{shape}(T[m]) - \text{shape}(S[m]).$$

On the other hand, Lemma 31 yields

$$\begin{aligned} 0 &\leq \text{shape}(\sigma_{\Gamma, i}(T)[x]) - \text{shape}(\sigma_{\Gamma', i}(S)[x]) = \\ &\text{shape}(T[x]) + \nu_x(\Gamma) - \text{shape}(S[x]) - \nu_x(\Gamma'), \end{aligned}$$

whence  $\text{shape}(T[x]) - \text{shape}(S[x]) \geq \nu_x(\Gamma') - \nu_x(\Gamma) > 0$ .  $\square$

**Corollary 36.** *Suppose that  $F = \sum_{S \in X} \alpha_S F_S$ , where  $X$  is a nonempty set of regular row standard  $\lambda$ -tableaux and  $\alpha_S \in \mathbf{K}^*$ . Let  $T$  be a minimal element of  $X$  and  $\Gamma$  be the smallest element of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ . If  $\sigma_{\Gamma, i}(T)$  is well-defined, then*

$$\xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F) = \sum_{P \in Y} \beta_P F_P,$$

where  $Y$  is a set of regular row standard  $(\lambda_1, \dots, \lambda_i, \lambda_{i+1}+1, \dots, \lambda_n+1)$ -tableaux,  $\beta_P \in \mathbf{K}^*$  and  $\sigma_{\Gamma, i}(T)$  is a minimal element of  $Y$ .

*Proof.* By Corollary 30, we obtain

$$\begin{aligned} \xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F) &= \sum \left\{ \alpha_S \text{sgn}_i(\Gamma') F_{\sigma_{\Gamma', i}(S)} \mid \right. \\ &\quad \left. \Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q), S \in X \text{ and } \sigma_{\Gamma', i}(S) \text{ is well-defined} \right\}. \end{aligned}$$

Therefore, it suffices to show the following fact: if  $\Gamma' \in \mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ ,  $S \in X$ ,  $\sigma_{\Gamma', i}(S)$  is well-defined and  $(\Gamma', S) \neq (\Gamma, T)$ , then  $\sigma_{\Gamma', i}(S) \not\leq \sigma_{\Gamma, i}(T)$ .

Indeed suppose that  $\sigma_{\Gamma', i}(S) \leq \sigma_{\Gamma, i}(T)$  for some  $S$  and  $\Gamma'$  satisfying the above hypothesis. Recall that  $\Gamma \leq \Gamma'$ , since  $\Gamma$  is the smallest. By Lemma 35,  $S < T$  if  $\Gamma < \Gamma'$ , which contradicts the minimality of  $T$ . If  $\Gamma = \Gamma'$  then Lemma 35 implies  $S \leq T$ . However, the case  $S = T$  is impossible since  $(\Gamma', S) \neq (\Gamma, T)$ . Hence, we again obtain the contradiction  $S < T$ .  $\square$

**3.7. Proof of Conjecture B.** Let  $\omega = c_1\omega_1 + \dots + c_{n-1}\omega_{n-1}$  be a nonzero dominant weight. Choose any  $i = 1, \dots, n-1$  such that  $c_i > 0$ . Consider sequences  $i_1^{(q)}, \dots, i_{k_q}^{(q)}; i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}; \dots; i_1^{(1)}, \dots, i_{k_1}^{(1)}$  and integers  $a_1, \dots, a_q, b_1, \dots, b_q$  satisfying the hypothesis of Lemma 27. For brevity, we shall use the abbreviation  $X_\alpha^V := X_{\alpha, 1}^V$ . By Lemma 10 and Definition 28, for any  $F \in \mathcal{U}^-$ , we have

$$\begin{aligned} &\left[ X_{\alpha_{i_1}^{(q)}}^V, \dots, X_{\alpha_{i_{k_q}}^{(q)}}^V, X_{\alpha_i}^V, X_{\alpha_{i_1}^{(q-1)}}^V, \dots, X_{\alpha_{i_{k_{q-1}}}^{(q-1)}}^V, X_{\alpha_i}^V, \dots, \right. \\ &\quad \left. X_{\alpha_{i_1}^{(1)}}^V, \dots, X_{\alpha_{i_{k_1}}^{(1)}}^V, X_{\alpha_i}^V, D_{\omega_i}^\omega \right] (F e_\omega^+) = \\ &\left[ \eta_{\alpha_{i_1}^{(q)}}, \dots, \eta_{\alpha_{i_{k_q}}^{(q)}}, \eta_{\alpha_i}, \eta_{\alpha_{i_1}^{(q-1)}}, \dots, \eta_{\alpha_{i_{k_{q-1}}}^{(q-1)}}, \eta_{\alpha_i}, \dots, \right. \\ &\quad \left. \eta_{\alpha_{i_1}^{(1)}}, \dots, \eta_{\alpha_{i_{k_1}}^{(1)}}, \eta_{\alpha_i}, \theta_{\omega_i} \right] (F) e_{\omega - \omega_i}^+ = \\ &\left[ i_1^{(q)}, \dots, i_{k_q}^{(q)}, i, i_1^{(q-1)}, \dots, i_{k_{q-1}}^{(q-1)}, i, \dots, i_1^{(1)}, \dots, i_{k_1}^{(1)}, i \right]_i (F) e_{\omega - \omega_i}^+ = \\ &\xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F) e_{\omega - \omega_i}^+, \end{aligned} \tag{21}$$

where  $V = \Delta(\omega) \oplus \Delta(\omega - \omega_i)$  and  $D_{\omega_i}^\omega$  is any operator on  $V$  whose restriction to  $\Delta(\omega)$  coincides with  $d_{\omega_i}^\omega$ . Hence the restriction

$$\left[ X_{\alpha_{i_1}^{(q)}}^V, \dots, X_{\alpha_{i_{k_q}}^{(q)}}^V, X_{\alpha_i}^V, X_{\alpha_{i_1}^{(q-1)}}^V, \dots, X_{\alpha_{i_{k_{q-1}}}^{(q-1)}}^V, X_{\alpha_i}^V, \dots, \right. \\ \left. X_{\alpha_{i_1}^{(1)}}^V, \dots, X_{\alpha_{i_{k_1}}^{(1)}}^V, X_{\alpha_i}^V, D_{\omega_i}^\omega \right] \Big|_{\Delta(\omega)} \quad (22)$$

is a map from  $\Delta(\omega)$  to  $\Delta(\omega - \omega_i)$  depending only on  $a_1, \dots, a_q, b_1, \dots, b_q, i$ . We denote it by  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)$ . Now we can rewrite (21) as

$$z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)(Fe_\omega^+) = \xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F)e_{\omega - \omega_i}^+. \quad (23)$$

**Lemma 37.** *Let  $\omega = c_1\omega_1 + \dots + c_{n-1}\omega_{n-1}$  be a nonzero dominant weight and  $v$  be a nonzero vector of  $\Delta(\omega)$ . Then there exist integers  $a_1, \dots, a_q, b_1, \dots, b_q, i$  such that*

$$1 \leq a_1 < a_2 < \dots < a_q \leq i < b_q < \dots < b_2 < b_1 \leq n, \quad c_i > 0$$

and  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)(v) \neq 0$ .

*Proof.* Let  $\lambda_j := \sum_{k=j}^{n-1} c_k$  for  $j = 1, \dots, n$ . Then  $\lambda := (\lambda_1, \dots, \lambda_n)$  is a partition coherent with  $\omega$ . Then by Proposition 29 we can write  $v = Fe_\omega^+$  with  $F = \sum_{S \in X} \alpha_S F_S$ , where  $X$  is a nonempty set consisting of standard  $\lambda$ -tableaux and  $\alpha_S \in \mathbf{K}^*$ . Let  $T$  be a minimal element of  $X$ . We denote by  $i$  the maximal number of  $1, \dots, n-1$  such that  $c_i > 0$ . Clearly,  $i$  is the height of the first column of  $[\lambda]$ .

Let  $i_0$  be the minimal number of  $1, \dots, i$  such that  $i_0 < T(i_0, 1)$  or be  $+\infty$  if there is no such number. Obviously,  $s = T(s, 1)$  for any  $s = 1, \dots, i_0 - 1$  and  $s < T(s, 1)$  for any  $s = i_0, \dots, i$ , since  $T$  is standard. Consider the graph  $\Gamma$  with vertices  $\mathbf{Z}$  and edges

$$(i, T(i, 1)), (i-1, T(i-1, 1)), \dots, (i_0, T(i_0, 1)).$$

The beginnings of all edges of  $\Gamma$  are mutually distinct and the ends of all edges of  $\Gamma$  are mutually distinct too. Therefore,  $\Gamma$  is a flow (see Definition 11).

Since the beginnings of the edges of  $\Gamma$  are less than or equal to  $i$ , the sources and transit points of  $\Gamma$  are also less than or equal to  $i$ . If  $(s, T(s, 1))$  is an edge of  $\Gamma$  and  $T(s, 1) \leq i$ , then  $i_0 < T(s, 1) \leq i$  and  $(T(s, 1), T(T(s, 1), 1))$  is an edge of  $\Gamma$  beginning at  $T(s, 1)$ . Hence  $T(s, 1)$  is a transit point of  $\Gamma$ . Therefore, all the sinks of  $\Gamma$  are greater than  $i$ .

We denote by  $a_1, \dots, a_q$  the sources of  $\Gamma$  and by  $b_1, \dots, b_q$  the sinks of  $\Gamma$  ordered so that (18) holds. Note that  $\Gamma$  can be the empty flow, in which case  $i_0 = +\infty$  and  $q = 0$ .

We are going to show that  $\Gamma$  is the smallest flow of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ . Take any flow  $\Gamma'$  of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$  distinct from  $\Gamma$ . In that case, we have  $q > 0$  and  $1 \leq i_0 \leq i$ . Let  $(s'_1, t'_1), \dots, (s'_{r'}, t'_{r'})$  be all the edges of  $\Gamma'$  ordered so that  $t'_1 > \dots > t'_{r'}$ . We set  $r := i - i_0 + 1$  and  $s_k := i - k + 1$ ,  $t_k := T(i - k + 1, 1)$  for any  $k := 1, \dots, r$ . Then the edges of  $\Gamma$  are  $(s_1, t_1), \dots, (s_r, t_r)$  and  $t_1 > \dots > t_r$ .

Following the comparison algorithm described in Section 3.6, we take  $l = 1, \dots, \min\{r, r'\}$  such that  $(s_k, t_k) = (s'_k, t'_k)$  for  $k < l$  and  $(s_l, t_l) \neq (s'_l, t'_l)$ . We have  $t'_l = t_l$  and  $s'_l \neq s_l$ . Moreover,  $s'_l \leq i$  and  $s'_l \notin \{s'_1, \dots, s'_{l-1}, s_l\} =$

$\{s_1, \dots, s_l\} = \{i, \dots, i-l+1\}$ , since  $\Gamma'$  is a flow of  $\mathcal{F}_i(a_1, \dots, a_q; b_1, \dots, b_q)$ . Hence  $s'_l < i-l+1 = s_l$  and  $\Gamma' > \Gamma$ .

By (23), we have

$$z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)(v) = \xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F)e_{\omega-\omega_i}^+.$$

It is easy to see that  $\sigma_{\Gamma,i}(T)$  is well-defined and is the tableau obtained from  $T$  by replacing its first column by the column of height  $n$  having  $k$  in each row  $k = 1, \dots, n$ . Clearly,  $\sigma_{\Gamma,i}(T)$  is standard.

By Corollary 36, we have

$$\xi_i(a_1, \dots, a_q; b_1, \dots, b_q)(F) = \sum_{P \in Y} \beta_P F_P,$$

where  $Y$  is a set of regular row standard  $(\lambda_1, \dots, \lambda_i, \lambda_{i+1}+1, \dots, \lambda_n+1)$ -tableaux,  $\beta_P \in \mathbf{K}^*$  and  $\sigma_{\Gamma,i}(T)$  is a minimal element of  $Y$ . Suppose that some  $P \in Y$  is not standard. Note that  $(\lambda_1, \dots, \lambda_i, \lambda_{i+1}+1, \dots, \lambda_n+1)$  is coherent with  $\omega - \omega_i$ . Then by Proposition 32,  $F_P e_{\omega-\omega_i}^+$  is a linear combination of vectors  $F_Q e_{\omega-\omega_i}^+$ , where  $Q$  is a standard  $(\lambda_1, \dots, \lambda_i, \lambda_{i+1}+1, \dots, \lambda_n+1)$ -tableau such that  $Q > P$ . In particular,  $Q \neq \sigma_{\Gamma,i}(T)$  by virtue of the minimality of  $\sigma_{\Gamma,i}(T)$  in  $Y$ . Hence the vector  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)(v)$  written in the standard basis of  $\Delta(\omega - \omega_i)$  has a nonzero coefficient at  $F_{\sigma_{\Gamma,i}(T)} e_{\omega-\omega_i}^+$  and thus is itself nonzero (Proposition 29).  $\square$

**Remark 38.** In this proof, it is important that all tableaux of  $Y$  including  $\sigma_{\Gamma,i}(T)$  have shape coherent with  $\omega - \omega_i$ . (Otherwise we would not be able to apply Propositions 29 and 32). We included step (ii) in the definition of  $\sigma_{\Gamma,i}(T)$  exactly to ensure this property.

**Theorem 39.** *If  $G = A_{n-1}(\mathbf{K})$ , then Conjecture B holds for all  $(F, \omega)$ .*

*Proof.* We are going to apply Lemma 8. Let  $\omega$  be a nonzero dominant weight and  $v$  be a simply primitive vector belonging to (5). We are going to apply the operator  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)$  to  $v$ , where  $\omega - \omega_i$  is dominant and (18) holds. Recall that  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)$  equals (22), which in turn is a linear combination with coefficients  $\pm 1$  or 0 of the restrictions  $w|_{\Delta(\omega)}$ , where  $w$  is a product of the operators

$$\begin{aligned} & X_{\alpha_{i_1^{(q)}}}^V, \dots, X_{\alpha_{i_{k_q}^{(q)}}}^V, X_{\alpha_i}^V, X_{\alpha_{i_1^{(q-1)}}}^V, \dots, X_{\alpha_{i_{k_{q-1}}}^{(q-1)}}^V, X_{\alpha_i}^V, \dots, \\ & X_{\alpha_{i_1^{(1)}}}^V, \dots, X_{\alpha_{i_{k_1}^{(1)}}}^V, X_{\alpha_i}^V, D_{\omega_i}^\omega \end{aligned}$$

taken in an arbitrary order. If the last factor of  $w$  is  $D_{\omega_i}^\omega$ , then  $w|_{\Delta(\omega)}(v) = 0$ , since  $v$  belongs to (5) and the restriction of  $D_{\omega_i}^\omega$  to  $\Delta(\omega)$  is  $d_{\omega_i}^\omega$ . Otherwise  $w|_{\Delta(\omega)}(v) = 0$ , since  $v$  is simply primitive. Thus we have proved that  $z_i^\omega(a_1, \dots, a_q; b_1, \dots, b_q)(v) = 0$ . Hence  $v = 0$  by Lemma 37.  $\square$

## 4. APPENDIX: LIST OF NOTATIONS

|                                                    |                                                                                                                                                        |
|----------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathbf{Z}^+$                                     | set of nonnegative integers;                                                                                                                           |
| $\mathcal{U}$                                      | hyperalgebra of algebraic group $G$ , p. 1;                                                                                                            |
| $X_{\alpha,m}, H_{\alpha,m}$                       | images of $X_{\alpha}^m/m!$ and $\binom{H_{\alpha}}{m}$ in $\mathcal{U}$ respectively;                                                                 |
| $\mathcal{U}^-$                                    | subalgebra of $\mathcal{U}$ generated by all $X_{\alpha,m}$ with $\alpha < 0$ ;                                                                        |
| $\mathcal{U}^0$                                    | subalgebra of $\mathcal{U}$ generated by all $H_{\alpha,m}$ ;                                                                                          |
| $\mathcal{U}^{-,0}$                                | $\mathcal{U}^- \mathcal{U}^0$ ;                                                                                                                        |
| $X_{\alpha,m}^V$                                   | operator on a rational $G$ -module $V$ acting as the left multiplication by $X_{\alpha,m}$ ;                                                           |
| $X(T), X^+(T)$                                     | sets of weights and dominant weights of torus $T$ , p. 2;                                                                                              |
| $L(\omega)$                                        | irreducible rational module with highest weight $\omega$ ;                                                                                             |
| $v_{\omega}^+$                                     | fixed nonzero vector of $L(\omega)$ having weight $\omega$ ;                                                                                           |
| $\Delta(\omega)$                                   | Weyl module with highest weight $\omega$ ;                                                                                                             |
| $e_{\omega}^+$                                     | fixed nonzero vector of $\Delta(\omega)$ having weight $\omega$ ;                                                                                      |
| $\nabla(\omega)$                                   | module contravariantly dual to $\Delta(\omega)$ ;                                                                                                      |
| $h(\omega)$                                        | $a_1 + \dots + a_{\ell}$ for $\omega = a_1\omega_1 + \dots + a_{\ell}\omega_{\ell}$ , p. 2;                                                            |
| $\text{ev}^{\omega}, \text{r}_{\alpha,m}^{\omega}$ | p. 3;                                                                                                                                                  |
| $d_{\delta}^{\omega}$                              | $\mathcal{U}^-$ -homomorphism $\Delta(\omega) \rightarrow \Delta(\omega - \delta)$ that takes $e_{\omega}^+$ to $e_{\omega-\delta}^+$ , Lemma 5, p. 4; |
| $\theta_{\delta}$                                  | p. 7;                                                                                                                                                  |
| $\eta_{\alpha,m}$                                  | p. 8;                                                                                                                                                  |
| $S_q$                                              | group of bijections of $\{1, \dots, q\}$ (symmetric group);                                                                                            |
| $\mathcal{F}_i(a_1, \dots, b_q)$                   | p. 10;                                                                                                                                                 |
| $\llbracket i_1, \dots, i_k \rrbracket_i$          | $[\eta_{\alpha_{i_1}}, \dots, \eta_{\alpha_{i_k}}, \theta_{\omega_i}]$ , p. 14;                                                                        |
| $\eta_{\alpha}$                                    | $\eta_{\alpha,1}$ ;                                                                                                                                    |
| $\delta_{\mathcal{P}}$                             | 1 if $\mathcal{P}$ is true and 0 otherwise;                                                                                                            |
| $H_l$                                              | $H_{\alpha_l,1}$ ;                                                                                                                                     |
| $\text{UT}(n)$                                     | set of $n \times n$ matrices with entries in $\mathbf{Z}$ having 0 on and under the main diagonal;                                                     |
| $N^s$                                              | $\sum_{b=1}^n N_{s,b}$ , the sum of elements in row $s$ of $N \in \text{UT}(n)$ ;                                                                      |
| $F^{(N)}$                                          | $\prod_{1 \leq a < b \leq n} E_{b,a}^{(N_{a,b})}$ , p. 15;                                                                                             |
| $\mathcal{O}(a, i, b)$                             | Definition 25, p. 17;                                                                                                                                  |
| $\xi_i(a_1, \dots, b_q)$                           | Definition 28, p. 21;                                                                                                                                  |
| $[\lambda]$                                        | diagram of a composition $\lambda$ , p. 21;                                                                                                            |
| $\text{Mat}_T$                                     | p. 21;                                                                                                                                                 |
| $F_T$                                              | $F^{(\text{Mat}_T)}$ ;                                                                                                                                 |
| $\varepsilon_i$                                    | $(0, \dots, 0, 1, 0, \dots, 0)$ of length $n$ with 1 at position $i$ ;                                                                                 |
| $\alpha(i, j)$                                     | $\varepsilon_i - \varepsilon_j$ ;                                                                                                                      |

|                               |                                                                                                      |
|-------------------------------|------------------------------------------------------------------------------------------------------|
| $\sigma_{\Gamma,i}(T)$        | p. 22;                                                                                               |
| $\text{shape}(Q)$             | $\mu$ if $Q$ is a $\mu$ -tableau, p. 23;                                                             |
| $T[m]$                        | tableau obtained from $T$ by removing all nodes with entry greater than $m$ , p. 23;                 |
| $\text{chain}(T)$             | $(\text{shape}(T[1]), \text{shape}(T[2]), \dots, \text{shape}(T[n])),$ p. 23;                        |
| $\nu_m(\Gamma)$               | $\sum \{\varepsilon_s \mid (s, t) \text{ is an edge of } \Gamma \text{ and } s \leq m < t\},$ p. 24; |
| $X_\alpha^V$                  | $X_{\alpha,1}^V;$                                                                                    |
| $z_i^\omega(a_1, \dots, b_q)$ | p. 28, eq. (23).                                                                                     |

# REFERENCES

- [1] A. Kleshchev J. Brundan and I. Suprunenko, Semisimple restrictions from  $\text{GL}(n)$  to  $\text{GL}(n-1)$ , *J. reine angew. Math.*, **500** (1998), 83–112.
- [2] J.C. Jantzen, Representations of algebraic groups, Mathematical surveys and monographs, vol. 107, *American Mathematical Society*, Providence, RI, 2003.
- [3] O. Mathieu, Filtrations of  $G$ -modules, *Ann. Sci. Ecole Norm. Sup. (4)*, **23** (1990), n. 4, 625–644.
- [4] R. Steinberg, Lectures on Chevalley Groups, New Haven, 1968.
- [5] R.W. Carter and G. Lusztig, On the modular representations of the general linear and symmetric groups, *Math. Z.*, **136** (1974), 193–242.
- [6] V. Shchigolev, Generalization of Modular Lowering Operators for  $\text{GL}_n$ , *Comm. Algebra* **36**, No. 4 (2008), 1250–1288.
- [7] R.L. Graham, D.E. Knuth, O. Patashnik, Concrete Mathematics, Second Edition, *Addison-Wesley*, 1998.
- [8] J. Humphreys, Introduction to Lie Algebras and Representation Theory, Graduate Texts in Math. 9, New-York etc., 1972 (Springer).

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS, LOMONOSOV MOSCOW STATE UNIVERSITY, LENINSKIYE GORY, MOSCOW, 119899, RUSSIA  
*E-mail address:* shchigolev.vladimir@yahoo.com